

Article

When Do Types Induce the Same Belief Hierarchy?

Andrés Perea¹ and Willemien Kets^{2,*}

¹ EpiCenter and Department of Quantitative Economics, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands; a.perea@maastrichtuniversity.nl

² MEDS, Kellogg School of Management, Northwestern University, Evanston, IL 60208-2001, USA

* Correspondence: willemien.kets@gmail.com

Academic Editor: Paul Weirich

Received: 22 August 2016 ; Accepted: 30 September 2016 ; Published: 9 October 2016

Abstract: Type structures are a simple device to describe higher-order beliefs. However, how can we check whether two types generate the same belief hierarchy? This paper generalizes the concept of a type morphism and shows that one type structure is contained in another if and only if the former can be mapped into the other using a generalized type morphism. Hence, every generalized type morphism is a hierarchy morphism and vice versa. Importantly, generalized type morphisms do not make reference to belief hierarchies. We use our results to characterize the conditions under which types generate the same belief hierarchy.

Keywords: types; belief hierarchies; epistemic game theory; morphisms

JEL Classification: C72

1. Introduction

Higher-order beliefs play a central role in game theory. Whether a player is willing to invest in a project, for example, may depend on what he or she thinks that his or her opponent thinks about the economic fundamentals, what he or she thinks that his or her opponent thinks that he or she thinks, and so on, up to arbitrarily high order (e.g., [1]). Higher-order beliefs can also affect economic conclusions in settings ranging from bargaining [2,3] and speculative trade [4] to mechanism design [5]. Higher-order beliefs about actions are central to epistemic characterizations, for example, of rationalizability [6,7], Nash equilibrium [8,9] and forward induction reasoning [10]. In principle, higher-order beliefs can be modeled explicitly, using belief hierarchies. For applications, the type structures introduced by Harsanyi [11] provide a simple, tractable modeling device to represent players' higher-order beliefs.

While type structures provide a convenient way to represent higher-order beliefs, it may be difficult to check whether types generate the same belief hierarchy. The literature has considered the following question: given two type structures, \mathcal{T} and \mathcal{T}' , is it the case that for every type in \mathcal{T} , there is a type in \mathcal{T}' that generates the same belief hierarchy? That is, is the type structure \mathcal{T} contained in \mathcal{T}' ?¹ The literature has considered two different tests to address this question, one based on hierarchy morphisms and one based on type morphisms. Hierarchy morphisms can be used to give a complete answer to this question: a type structure \mathcal{T} is contained in \mathcal{T}' if and only if there is a hierarchy morphism from the former to the latter. A problem with this test is that hierarchy morphisms make reference to belief hierarchies, as we shall see. Therefore, this test requires us to go outside the purview of type structures. The second test uses type morphisms. Type morphisms are defined solely in terms

¹ We follow the terminology of Friedenber and Meier [12] here. A stronger condition for \mathcal{T} to be contained in \mathcal{T}' is that \mathcal{T} can be embedded (using a type morphism) into \mathcal{T}' as a belief-closed subset [13]. Our results can be used to characterize conditions under which \mathcal{T} is contained in \mathcal{T}' in this stronger sense in a straightforward way.

of the properties of the type structures. However, the test based on type morphisms only provides a sufficient condition: if there is a type morphism from \mathcal{T} to \mathcal{T}' , then \mathcal{T} is contained in \mathcal{T}' [14]. However, as shown by Friedenber and Meier [12], the condition is not necessary: it may be that \mathcal{T}' contains \mathcal{T} , yet there is no type morphism from \mathcal{T} to \mathcal{T}' . The work in [12] also provides a range of conditions under which the condition is both necessary and sufficient. However, they do not directly address the question of whether there might be an alternate test (which provides conditions that are both necessary and sufficient) that does not require us to describe the belief hierarchies explicitly.

This paper provides such a test, by generalizing the notion of a type morphism. We show that a type structure is contained in another if and only if there is a generalized type morphism from the former to the latter. Therefore, a generalized type morphism is a hierarchy morphism and vice versa. Unlike the definition of hierarchy morphisms, the definition of generalized type morphisms does not make reference to belief hierarchies. Therefore, this test can be carried out without leaving the purview of type structures. Using this result, it is straightforward to verify whether two types generate the same belief hierarchy, as we show.

Hierarchy morphisms are used in a number of different settings. For example, they can be used to check whether types have the same rationalizable actions [15] and play an important role in the literature on the robustness to misspecifying the parameter set more generally; see, e.g., Ely and Peski [16] and Liu [17]. Hierarchy morphisms are also used to study the robustness of Bayesian-Nash equilibria to misspecifications of players' belief hierarchies [18,19] and in epistemic game theory. The current results make it possible to study these issues without describing players' belief hierarchies explicitly, using that every hierarchy morphism is a generalized type morphism and conversely.

A critical ingredient in the definition of a generalized type morphism is the σ -algebra on a player's type set, which separates his or her types if and only if they differ in the belief hierarchy that they generate. Mertens and Zamir ([13], p. 6) use this σ -algebra to define non-redundant type structures, and this σ -algebra also plays an important role in the work of Friedenber and Meier [12], where it is used to characterize the conditions under which hierarchy morphisms and type morphisms coincide. The work in [13] provides a nonconstructive definition of this σ -algebra, and [12] show that the σ -algebra defined by [13] is the σ -algebra generated by the functions that map types into belief hierarchies. We provide a constructive definition of this σ -algebra, by means of a type partitioning procedure that does not make reference to belief hierarchies.

While many of the ingredients that underlie our results are known in some form or another, we view the contribution of this paper as combining these ideas in a new way to generalize the concept of a type morphism, so that it provides a necessary and sufficient condition for a type structure to be contained in another that does not refer to belief hierarchies.

A number of papers has shown that the measurable structure associated with type structures can impose restrictions on reasoning [12,20–23]. This paper contributes to that literature in two ways. First, we elucidate the connection by constructing the measurable structure on type sets that is generated by players' higher-order beliefs. Second, we provide tools to easily go from the domain of type structures to the domain of belief hierarchies and vice versa.

The outline of this paper is as follows. The next section introduces basic concepts. Section 3 discusses type morphisms and hierarchy morphisms. Section 4 defines our generalization of a type morphism and proves the main result. Section 5 applies this result to characterize the conditions under which types generate the same belief hierarchy. Section 6 considers the special case where players have finitely many types. Proofs are relegated to the Appendix A.

2. Belief Hierarchies and Types

In this section, we show how belief hierarchies can be encoded by means of a type structure. The original idea behind this construction goes back to Harsanyi (1967). We first provide the definition of a type structure and subsequently explain how to derive a belief hierarchy from a type in a type structure. We conclude the section with an example of two type structures that are equivalent,

in the sense that they produce exactly the same sets of belief hierarchies for the players. This example thus shows that the same belief hierarchy can be encoded within different type structures.

2.1. Type Structures

Consider a finite set of players I . Assume that each player i faces a basic space of uncertainty (X_i, Σ_i) , where X_i is a set and Σ_i a σ -algebra on X_i . That is, $\mathcal{X}_i = (X_i, \Sigma_i)$ is a measurable space. The combination $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$ of basic uncertainty spaces is called a multi-agent uncertainty space. The basic space of uncertainty for player i could, for instance, be the set of opponents' choice combinations, or the set of parameters determining the utility functions of the players, or even a combination of the two.

A belief hierarchy for player i specifies a probability measure on \mathcal{X}_i , the first-order belief, a probability measure on \mathcal{X}_i and the opponents' possible first-order beliefs, the second-order belief, and so on. As is standard, we encode such infinite belief hierarchies by means of type structures.

For any measurable space $(Y, \hat{\Sigma})$, we denote by $\Delta(Y, \hat{\Sigma})$ the set of probability measures on $(Y, \hat{\Sigma})$. We endow $\Delta(Y, \hat{\Sigma})$ with the coarsest σ -algebra that contains the sets:

$$\{\mu \in \Delta(Y, \hat{\Sigma}) \mid \mu(E) \geq p\} : E \in \hat{\Sigma}, p \in [0, 1].$$

This is the σ -algebra used in Heifetz and Samet [14] and many subsequent papers; it coincides with the Borel σ -algebra on $\Delta(Y, \hat{\Sigma})$ (induced by the weak convergence topology) if Y is metrizable and $\hat{\Sigma}$ is the Borel σ -algebra. Product spaces are endowed with the product σ -algebra. Given a collection of measurable spaces (Y_i, \mathcal{Y}_i) , $i \in I$, write \mathcal{Y} for the product σ -algebra $\otimes_{j \in I} \mathcal{Y}_j$ and \mathcal{Y}_{-i} for the product σ -algebra $\otimes_{j \neq i} \mathcal{Y}_j$, where $i \in I$.

Definition 1. (Type structure) Consider a multi-agent uncertainty space $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$. A type structure for \mathcal{X} is a tuple $\mathcal{T} = (T_i, \Sigma_i^{\mathcal{T}}, b_i)_{i \in I}$ where, for every player i ,

- (a) T_i is a set of types for player i , endowed with a σ -algebra $\Sigma_i^{\mathcal{T}}$, and
- (b) $b_i : T_i \rightarrow \Delta(X_i \times T_{-i}, \hat{\Sigma}_i)$ is a measurable mapping that assigns to every type t_i a probabilistic belief $b_i(t_i) \in \Delta(X_i \times T_{-i}, \hat{\Sigma}_i)$ on its basic uncertainty space and the opponents' type combinations, where $\hat{\Sigma}_i = \Sigma_i \otimes \Sigma_{-i}^{\mathcal{T}}$ is the product σ -algebra on $X_i \times T_{-i}$.

Finally, if $f : Y \rightarrow (Y', \Sigma')$ is a function from Y to the measurable space (Y', Σ') , then $\sigma(f)$ is the σ -algebra on Y generated by f , that is, it is the coarsest σ -algebra that contains the sets $\{y \in Y : f(y) \in E\}$ for $E \in \Sigma'$.

2.2. From Type Structures to Belief Hierarchies

In the previous subsection, we have introduced the formal definition of a type structure. We now show how to "decode" a type within a type structure, by deriving the full belief hierarchy that it induces.

Consider a type structure $\mathcal{T} = (T_i, \Sigma_i^{\mathcal{T}}, b_i)_{i \in I}$ for \mathcal{X} . Then, every type t_i within \mathcal{T} induces an infinite belief hierarchy:

$$h_i^{\mathcal{T}}(t_i) = (\mu_i^{\mathcal{T},1}(t_i), \mu_i^{\mathcal{T},2}(t_i), \dots),$$

where $\mu_i^{\mathcal{T},1}(t_i)$ is the induced first-order belief, $\mu_i^{\mathcal{T},2}(t_i)$ is the induced second-order belief, and so on. We will inductively define, for every n , the n -th order beliefs induced by types t_i in \mathcal{T} , building upon the $(n - 1)$ -th order beliefs that have been defined in the preceding step.

We start by defining the first-order beliefs. For each player i , define:

$$H_i^1 := \Delta(X_i, \Sigma_i)$$

to be the set of beliefs about X_i , and for every type $t_i \in T_i$, define its first-order belief $\mu_i^{\mathcal{T},1}(t_i)$ by:

$$\mu_i^{\mathcal{T},1}(t_i)(E_i) := b_i(t_i)(E_i \times T_{-i}) \text{ for all } E_i \in \Sigma_i.$$

Clearly, $\mu_i^{\mathcal{T},1}(t_i) \in \Delta(X_i, \Sigma_i)$ for every type t_i . Define $h_i^{\mathcal{T},1}(t_i) := \mu_i^{\mathcal{T},1}(t_i)$. The mapping $\mu_i^{\mathcal{T},1}$ from T_i to H_i^1 is measurable by standard arguments. For $n > 1$, suppose the set H_i^{n-1} has been defined and that the function $h_i^{\mathcal{T},n-1}$ from T_i to H_i^{n-1} is measurable. Let $\hat{\Sigma}_i^{n-1}$ be the product σ -algebra on $X_i \times \times_{j \neq i} H_j^{n-1}$, and define:

$$H_i^n := H_i^{n-1} \times \Delta(X_i \times H_{-i}^{n-1}, \hat{\Sigma}_i^{n-1}).$$

For every type t_i , define its n -th-order belief $\mu_i^{\mathcal{T},n}(t_i)$ by:

$$\text{for all } E \in \hat{\Sigma}_i^{n-1} : \quad \mu_i^{\mathcal{T},n}(t_i)(E) = b_i(t_i)(\{(x_i, t_{-i}) \in X_i \times T_{-i} \mid (x_i, h_{-i}^{\mathcal{T},n-1}(t_{-i})) \in E\}),$$

with $h_{-i}^{\mathcal{T},n-1}(t_{-i}) = (h_j^{\mathcal{T},n-1}(t_j))_{j \neq i}$. Since $h_j^{\mathcal{T},n-1}$ is measurable for every player j , $\mu_i^{\mathcal{T},n}$ is indeed a probability measure on $(X_i \times H_{-i}^{n-1}, \hat{\Sigma}_i^{n-1})$. Define $h_i^{\mathcal{T},n}(t_i) := (h_i^{\mathcal{T},n-1}(t_i), \mu_i^{\mathcal{T},n}(t_i))$. It follows that $h_i^{\mathcal{T},n}(t_i) \in H_i^n$. Moreover, $h_i^{\mathcal{T},n}$ is measurable.

Note that, formally speaking, the n -th-order belief $\mu_i^{\mathcal{T},n}(t_i)$ is a belief about X_i and the opponents' first-order until $(n - 1)$ -th order beliefs. Moreover, by construction, the n -th and $(n + 1)$ -th order beliefs $\mu_i^{\mathcal{T},n}(t_i)$ and $\mu_i^{\mathcal{T},n+1}(t_i)$ are coherent in the sense that they induce the same belief on X_i and the opponents' first-order until $(n - 1)$ -th order beliefs.

Finally, for every type $t_i \in T_i$, we denote by:

$$h_i^{\mathcal{T}}(t_i) := (\mu_i^{\mathcal{T},n}(t_i))_{n \in \mathbb{N}}$$

the belief hierarchy induced by type t_i in \mathcal{T} . Furthermore, define H_i to be the set $\Delta(X_i) \times \times_{n \geq 1} \Delta(X_i \times H_{-i}^n)$ of all belief hierarchies. We say that two types, t_i and t'_i , of player i generate the same belief hierarchy if $h_i^{\mathcal{T}}(t_i) = h_i^{\mathcal{T}}(t'_i)$. Types t_i and t'_i generate the same n -th-order belief if $\mu_i^{\mathcal{T},n}(t_i) = \mu_i^{\mathcal{T},n}(t'_i)$.²

2.3. Example

Consider a multi-agent uncertainty space $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$ where $I = \{1, 2\}$, $X_1 = \{c, d\}$, $X_2 = \{e, f\}$ and Σ_1, Σ_2 are the discrete σ -algebras on X_1 and X_2 , respectively. Consider the type structures $\mathcal{T} = (T_1, T_2, \Sigma_1, \Sigma_2, b_1, b_2)$ and $\mathcal{T}' = (R_1, R_2, \Sigma_1, \Sigma_2, \beta_1, \beta_2)$ in Table 1.

² Clearly, t_i and t'_i generate the same n -th-order belief if and only if $h_i^{\mathcal{T},n}(t_i) = h_i^{\mathcal{T},n}(t'_i)$.

Table 1. Two equivalent type structures.

Type Structure \mathcal{T}
$T_1 = \{t_1, t'_1, t''_1\}, \quad T_2 = \{t_2, t'_2, t''_2\}$
$b_1(t_1) = \frac{1}{2}(c, t_2) + \frac{1}{2}(d, t'_2)$ $b_1(t'_1) = \frac{1}{6}(c, t_2) + \frac{1}{3}(c, t'_2) + \frac{1}{2}(d, t'_2)$ $b_1(t''_1) = \frac{1}{2}(c, t'_2) + \frac{1}{2}(d, t'_2)$
$b_2(t_2) = \frac{1}{4}(e, t_1) + \frac{1}{2}(e, t'_1) + \frac{1}{4}(f, t''_1)$ $b_2(t'_2) = \frac{1}{8}(e, t_1) + \frac{1}{8}(e, t'_1) + \frac{3}{4}(f, t''_1)$ $b_2(t''_2) = \frac{3}{8}(e, t_1) + \frac{3}{8}(e, t'_1) + \frac{1}{4}(f, t''_1)$
Type Structure \mathcal{T}'
$R_1 = \{r_1, r'_1, r''_1\}, \quad R_2 = \{r_2, r'_2, r''_2\}$
$\beta_1(r_1) = \frac{1}{4}(c, r_2) + \frac{1}{4}(c, r'_2) + \frac{1}{2}(d, r'_2)$ $\beta_1(r'_1) = \frac{1}{2}(c, r'_2) + \frac{1}{8}(d, r_2) + \frac{3}{8}(d, r'_2)$ $\beta_1(r''_1) = \frac{1}{2}(c, r'_2) + \frac{3}{8}(d, r_2) + \frac{1}{8}(d, r'_2)$
$\beta_2(r_2) = \frac{1}{4}(e, r'_1) + \frac{3}{4}(f, r_1)$ $\beta_2(r'_2) = \frac{3}{4}(e, r'_1) + \frac{1}{4}(f, r_1)$ $\beta_2(r''_2) = \frac{1}{8}(e, r'_1) + \frac{1}{8}(e, r''_1) + \frac{3}{4}(f, r_1)$

Then, it can be verified that the types t_1, t'_1, r'_1 and r''_1 generate the same belief hierarchy, and so do the types t'_1 and r_1 , the types t_2, t'_2 and r'_2 and the types t'_2, r_2 and r''_2 . In particular, for every type in \mathcal{T} , there is another type in \mathcal{T}' generating the same belief hierarchy, and vice versa. In this sense, the two type structures \mathcal{T} and \mathcal{T}' are equivalent.

3. Hierarchy and Type Morphisms

The literature has considered two concepts that map type structures into each other, type morphisms and hierarchy morphisms. Throughout the remainder of the paper, fix two type structures, $\mathcal{T} = (T_i, \Sigma_i^{\mathcal{T}}, b_i)_{i \in I}$ and $\mathcal{T}' = (T'_i, \Sigma_i^{\mathcal{T}'}, b'_i)_{i \in I}$ on \mathcal{X} . The functions that map types from \mathcal{T} and \mathcal{T}' into belief hierarchies are denoted by $h_i^{\mathcal{T}}$ and $h_i^{\mathcal{T}'}$, respectively.

Definition 2. (Hierarchy morphism) For each player $i \in I$, let φ_i be a function from T_i to T'_i , such that for every type $t_i \in T_i$, $h_i^{\mathcal{T}'}(\varphi_i(t_i)) = h_i^{\mathcal{T}}(t_i)$. Then, φ_i is a hierarchy morphism (from \mathcal{T} to \mathcal{T}'). With some abuse of notation, we refer to the profile $(\varphi_i)_{i \in I}$ as a hierarchy morphism.

Therefore, if there is a hierarchy morphism between \mathcal{T} and \mathcal{T}' , then every type in \mathcal{T} can be mapped into a type in \mathcal{T}' in a way that preserves belief hierarchies. We say that the type structure \mathcal{T}' contains \mathcal{T} if, and only if, there is a hierarchy morphism from \mathcal{T} to \mathcal{T}' .

Type morphisms are mappings between type structures that preserve beliefs.

Definition 3. (Type morphism) For each player $i \in I$, let φ_i be a function from T_i to T'_i that is measurable with respect to $\Sigma_i^{\mathcal{T}}$ and $\Sigma_i^{\mathcal{T}'}$.³ Suppose that for each player i , type $t_i \in T_i$ and $E \in \Sigma_i \otimes \Sigma_{-i}^{\mathcal{T}'}$,

$$b_i(t_i)(\{(x_i, t_{-i}) \in X_i \times T_{-i} \mid (x_i, \varphi_{-i}(t_{-i})) \in E\}) = b'_i(\varphi_i(t_i))(E). \tag{1}$$

Then, $\varphi := (\varphi_i)_{i \in I}$ is a type morphism (from \mathcal{T} to \mathcal{T}').

³ That is, for each $E \in \Sigma_i^{\mathcal{T}'}$, we have $\{t_i \in T_i \mid \varphi_i(t_i) \in E\} \in \Sigma_i^{\mathcal{T}}$.

Heifetz and Samet [14] have shown that one type structure is contained in another whenever there is a type morphism from the former to the latter.

Proposition 1. ([14], Prop. 5.1) *If φ is a type morphism from \mathcal{T} to \mathcal{T}' , then it is a hierarchy morphism. Therefore, if there is a type morphism from \mathcal{T} to \mathcal{T}' , then \mathcal{T}' contains \mathcal{T} .*

Unlike hierarchy morphisms, type morphisms do not make reference to belief hierarchies. Therefore, to check whether there is a type morphism from one type structure to another, we need to consider only the type structures. However, the condition that there be a type morphism from one type structure to another provides only a sufficient condition for the former to be contained in the latter. Indeed, Friedenberg and Meier [12] show that the condition is not necessary: there are type structures such that one is contained in the other, yet there is no type morphism between the two.

4. Generalized Type Morphisms

Type morphisms require beliefs to be preserved for every event in the types' σ -algebra. However, for two types to generate the same belief hierarchy, it suffices that their beliefs are preserved only for events that can be described in terms of players' belief hierarchies. We use this insight to define generalized type morphisms and show that a type structure contains another if and only if there is a generalized type morphism from the latter to the former.

The first step is to define the relevant σ -algebra. Mertens and Zamir ([13], p. 6) provide the relevant condition. We follow the presentation of Friedenberg and Meier [12].

Definition 4. ([12], Def. 5.1) *Fix a type structure \mathcal{T} and fix a sub- σ algebra $\tilde{\Sigma}_i^{\mathcal{T}} \subseteq \Sigma_i^{\mathcal{T}}$ for each player $i \in I$. Then, the product σ -algebra $\tilde{\Sigma}^{\mathcal{T}}$ is closed under \mathcal{T} if for each player i ,*

$$\{t_i \in T_i \mid b_i(t_i)(E) \geq p\} \in \tilde{\Sigma}_i^{\mathcal{T}}$$

for all $E \in \Sigma_i \otimes \tilde{\Sigma}_{-i}^{\mathcal{T}}$ and $p \in [0, 1]$.

The coarsest (sub-) σ algebra that is closed under \mathcal{T} is of special interest, and we denote it by $\mathcal{F}^{\mathcal{T}} = \bigotimes_{i \in I} \mathcal{F}_i^{\mathcal{T}}$. It is the intersection of all σ -algebras that are closed under \mathcal{T} .⁴ The work in [13] uses this σ -algebra to define non-redundant type spaces, and [12] use it to characterize the condition under which a hierarchy morphism is a type morphism.

Friedenberg and Meier [12] provide a characterization of the σ -algebra $\mathcal{F}^{\mathcal{T}}$ in terms of the hierarchy mappings. Recall that $\sigma(h_i^{\mathcal{T}})$ is the σ -algebra on T_i generated by the mapping $h_i^{\mathcal{T}}$. That is, $\sigma(h_i^{\mathcal{T}})$ is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid h_i^{\mathcal{T}}(t_i) \in E\} : \quad E \subseteq H_i \text{ measurable.}$$

Lemma 1. ([12], Lemma 6.4) *Let the product σ -algebra $\mathcal{F}^{\mathcal{T}}$ be the coarsest σ -algebra that is closed under \mathcal{T} . Then, for each player i , $\mathcal{F}_i^{\mathcal{T}} = \sigma(h_i^{\mathcal{T}})$.*

We are now ready to define generalized type morphisms.

⁴ Since $\Sigma^{\mathcal{T}}$ is closed under \mathcal{T} (by measurability of the belief maps b_i), the intersection is nonempty. It is easy to verify that the intersection is a σ -algebra.

Definition 5. (Generalized type morphism) For each player $i \in I$, let φ_i be a function from T_i to T'_i that is measurable with respect to $\Sigma_i^{\mathcal{T}}$ and $\mathcal{F}_i^{\mathcal{T}'}$.⁵ Suppose that for each player i , type $t_i \in T_i$ and $E \in \Sigma_i \otimes \mathcal{F}_{-i}^{\mathcal{T}'}$,

$$b_i(t_i)(\{(x_i, t_{-i}) \in X_i \times T_{-i} \mid (x_i, \varphi_{-i}(t_{-i})) \in E\}) = b'_i(\varphi_i(t_i))(E).$$

Then, $\varphi := (\varphi_i)_{i \in I}$ is a generalized type morphism (from \mathcal{T} to \mathcal{T}').

Note that a type morphism is always a generalized type morphism, but not vice versa. Like type morphisms, generalized type morphisms are defined using the language of type structures alone; the definition does not make reference to belief hierarchies. The difference between type morphisms and generalized type morphisms is that the former requires beliefs to be preserved for all events in the σ -algebra $\Sigma_i \otimes \Sigma_{-i}^{\mathcal{T}'}$ for player i , while the latter requires beliefs to be preserved only for events in the σ -algebra $\Sigma_i \otimes \mathcal{F}_{-i}^{\mathcal{T}'}$, and this σ -algebra is a coarsening of $\Sigma_i \otimes \Sigma_{-i}^{\mathcal{T}'}$ (Definition 4 and Lemma 1).

Our main result states that one structure is contained in another if and only if there is a generalized type morphism from the former to the latter.

Theorem 1. A mapping φ is a hierarchy morphism from \mathcal{T} to \mathcal{T}' if and only if it is a generalized type morphism from \mathcal{T} to \mathcal{T}' . Hence, a type structure \mathcal{T}' contains \mathcal{T} if and only if there is a generalized type morphism from \mathcal{T} to \mathcal{T}' .

This result establishes an equivalence between generalized type morphisms and hierarchy morphisms. It thus provides a test that can be used to verify whether one type structure is contained in the other that does not refer to belief hierarchies.

While the characterization in Theorem 1 does not make reference to belief hierarchies, the result may not be easy to apply directly. The σ -algebras $\mathcal{F}_i^{\mathcal{T}}$ are defined as the intersection of σ -algebras that are closed under \mathcal{T} , and there can be (uncountably) many of those. We next define a simple procedure to construct this σ -algebra.

Procedure 1. (Type partitioning procedure) Consider a multi-agent uncertainty space $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$ and a type structure $\mathcal{T} = (T_i, \Sigma_i^{\mathcal{T}}, b_i)_{i \in I}$ for \mathcal{X} .

Initial step: For every player i , let $\mathcal{S}_i^{\mathcal{T},0} = \{T_i, \emptyset\}$ be the trivial σ -algebra of his or her set of types T_i .

Inductive step: Suppose that $n \geq 1$ and that the sub- σ algebra $\mathcal{S}_i^{\mathcal{T},n-1}$ on T_i has been defined for every player i . Then, for every player i , let $\mathcal{S}_i^{\mathcal{T},n}$ be the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid b_i(t_i)(E) \geq p\}$$

for all $E \in \Sigma_i \otimes \mathcal{S}_{-i}^{\mathcal{T},n-1}$ and all $p \in [0, 1]$. Furthermore, let $\mathcal{S}_i^{\mathcal{T},\infty}$ be the σ -algebra generated by the union $\bigcup_n \mathcal{S}_i^{\mathcal{T},n}$.

A simple inductive argument shows that $\mathcal{S}_i^{\mathcal{T},n}$ refines $\mathcal{S}_i^{\mathcal{T},n-1}$ for all players i and all n ; clearly, $\mathcal{S}_i^{\mathcal{T},\infty}$ refines $\mathcal{S}_i^{\mathcal{T},n}$ for any n . The next result shows that the type partitioning procedure delivers the σ -algebras that are generated by the hierarchy mappings.

Proposition 2. Fix a type structure \mathcal{T} , and let $i \in I$. Then, $\mathcal{S}_i^{\mathcal{T},\infty} = \sigma(h_i^{\mathcal{T}})$ and $\mathcal{S}_i^{\mathcal{T},n} = \sigma(h_i^{\mathcal{T},n})$ for all $n \geq 1$. Therefore, $\mathcal{S}_i^{\mathcal{T},\infty} = \mathcal{F}_i^{\mathcal{T}}$.

⁵ That is, for each $E \in \mathcal{F}_i^{\mathcal{T}'}$, we have $\{t_i \in T_i \mid \varphi_i(t_i) \in E\} \in \Sigma_i^{\mathcal{T}}$.

Hence, we can use the type partitioning procedure to construct the σ -algebras, which we need for our characterization result (Theorem 1). Heifetz and Samet [24] consider a similar procedure in the context of knowledge spaces to show that a universal space does not exist for that setting. The procedure also has connections with the construction in Kets [22] of type structures that describe the beliefs of players with a finite depth of reasoning. In the next section, we use Theorem 1 and the type partitioning procedure to characterize the types that generate the same belief hierarchies.

5. Characterizing Types with the Same Belief Hierarchy

We can use the results in the previous section to provide simple tests to determine whether two types, from the same type structure or from different structures, generate the same belief hierarchy. We assume in this section that \mathcal{X}_i is countably generated: there is a countable collection of subsets $E_i^n \subseteq X_i, n = 1, 2, \dots$, such that Σ_i is the coarsest σ -algebra that contains these subsets. Examples of countably-generated σ -algebras include the discrete σ -algebra on a finite or countable set and the Borel σ -algebra on a finite-dimensional Euclidean space. Recall that an atom of a σ -algebra Σ on a set Y is a set $a \in \Sigma$, such that Σ does not contain a nonempty proper subset of a . That is, for any $a' \in \Sigma$, such that $a' \subseteq a$, we have $a' = a$ or $a' = \emptyset$.⁶

Lemma 2. *Let $i \in I$ and $n \geq 1$. The σ -algebras $\mathcal{S}_i^{\mathcal{T},n}$ and $\mathcal{S}_i^{\mathcal{T},\infty}$ are atomic. That is, for each $t_i \in T_i$, there are atoms $a_i^n(t_i)$ and $a_i^\infty(t_i)$ in $\mathcal{S}_i^{\mathcal{T},n}$ and $\mathcal{S}_i^{\mathcal{T},\infty}$, respectively, such that $t_i \in a_i^n(t_i)$ and $t_i \in a_i^\infty(t_i)$.*

This result motivates the name “type partitioning procedure”: the procedure constructs a σ -algebra that partitions the type sets into atoms. Proposition 3 shows that these atoms contain precisely the types that generate the same higher-order beliefs.

Proposition 3. *For every player i , every $n \geq 1$ and every two types $t_i, t'_i \in T_i$, we have that*

(a) *for every $n \geq 0$, types t_i and t'_i generate the same n -th-order belief if and only if there is an atom $a_i^n \in \mathcal{S}_i^{\mathcal{T},n}$, such that $t_i, t'_i \in a_i^n$;*

(b) *types t_i and t'_i generate the same belief hierarchy if and only if there is an atom $a_i^\infty \in \mathcal{S}_i^{\mathcal{T},\infty}$, such that $t_i, t'_i \in a_i^\infty$.*

There is a connection between Proposition 3 and the work of Mertens and Zamir [13]. The work in [13] defines a type structure \mathcal{T} to be non-redundant if for every player i , the σ -algebra $\mathcal{F}_i^{\mathcal{T}}$ separates types; see Liu ([17], Prop. 2) for a result that shows that this definition is equivalent to the requirement that there are no two types that generate the same belief hierarchy. Therefore, [13] already note the connection between the separating properties of $\mathcal{F}^{\mathcal{T}}$ and the question of whether types generate the same belief hierarchy. The contribution of Proposition 3 is to provide a simple procedure to construct the σ -algebra $\mathcal{F}^{\mathcal{T}}$ and to show that the separating sets can be taken to be atoms (as long as the σ -algebra on X_i is countably generated).

Proposition 3 can also be used to verify whether two types from different type structures generate the same higher-order beliefs, by merging the two structures. Specifically, consider two different type structures, $\mathcal{T}^1 = (T_i^1, \Sigma_i^1, b_i^1)_{i \in I}$ and $\mathcal{T}^2 = (T_i^2, \Sigma_i^2, b_i^2)_{i \in I}$, for the same multi-agent uncertainty space $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$. To check whether two types $t_i^1 \in T_i^1$ and $t_i^2 \in T_i^2$ induce the same belief hierarchy, we can merge the two type structures into one large type structure and then run the type partitioning procedure on this larger type structure. That is, define the type structure $\mathcal{T}^* = (T_i^*, \Sigma_i^*, b_i^*)_{i \in I}$ as follows.

⁶ Clearly, for any $y \in Y$, if there is an atom a that contains y (i.e., $y \in a$), then this atom is unique.

For each player i , let T_i^* be the union of T_i^1 and T_i^2 (possibly made disjoint by replacing T_i^1 or T_i^2 with a homeomorphic copy), and define the σ -algebra Σ_i^* on T_i^* by:

$$E \in \Sigma_i^* \text{ if and only if } E \cap T_i^1 \in \Sigma_i^1 \text{ and } E \cap T_i^2 \in \Sigma_i^2.$$

Furthermore, define b_i^* by:

$$b_i^*(t_i) := \begin{cases} b_i^1(t_i), & \text{if } t_i \in T_i^1 \\ b_i^2(t_i), & \text{if } t_i \in T_i^2 \end{cases}$$

for all types $t_i \in T_i$.⁷ Applying the type partitioning procedure on \mathcal{T}^* gives a σ -algebra $\mathcal{S}_i^{*,\infty}$ on T_i^* for each player i . If $t_i^1 \in T_i^1$ and $t_i^2 \in T_i^2$ belong to the same atom of $\mathcal{S}_i^{*,\infty}$, then t_i^1 and t_i^2 induce the same belief hierarchy. The converse also holds, and hence, we obtain the following result.

Proposition 4. Consider two type structures $T^1 = (T_i^1, \Sigma_i^1, b_i^1)_{i \in I}$ and $T^2 = (T_i^2, \Sigma_i^2, b_i^2)_{i \in I}$. Let $T^* = (T_i^*, \Sigma_i^*, b_i^*)_{i \in I}$ be the large type structure defined above, obtained by merging the two type structures, and let $\mathcal{S}_i^{*,\infty}$, for a given player i , be the σ -algebra on T_i^* generated by the type partitioning procedure. Then, two types $t_i^1 \in T_i^1$ and $t_i^2 \in T_i^2$ induce the same belief hierarchy, if and only if, t_i^1 and t_i^2 belong to the same atom of $\mathcal{S}_i^{*,\infty}$.

The type partitioning procedure is thus an easy and effective way to check whether two types, from possibly different type structures, generate the same belief hierarchy or not.

We expect our main results to apply more broadly. The proofs can easily be modified so that the main results extend to conditional probability systems in dynamic games [25], lexicographic beliefs [26], beliefs of players with a finite depth of reasoning [22,27] and the Δ -hierarchies introduced by Ely and Peski [16].

6. Finite Type Structures

When type structures are finite, our results take on a particularly simple and intuitive form. Say that a type structure \mathcal{T} is finite if the type set T_i is finite for every player i . For finite type structures, we can replace σ -algebras by partitions.

We first define the type partitioning procedure for the case of finite type structures. A finite partition of a set A is a finite collection $\mathcal{P} = \{P_1, \dots, P_K\}$ of nonempty subsets $P_k \subseteq A$, such that $\bigcup_{k=1}^K P_k = A$ and $P_k \cap P_m = \emptyset$ whenever $k \neq m$. We refer to the sets P_k as equivalence classes. For an element $a \in A$, we denote by $\mathcal{P}(a)$ the equivalence class P_k to which a belongs. The trivial partition of A is the partition $\mathcal{P} = \{A\}$ containing a single set; the full set A . For two partitions \mathcal{P}^1 and \mathcal{P}^2 on A , we say that \mathcal{P}^1 is a refinement of \mathcal{P}^2 if for every set $P^1 \in \mathcal{P}^1$, there is a set $P^2 \in \mathcal{P}^2$, such that $P^1 \subseteq P^2$.

In the procedure, we recursively partition the set of types of an agent into equivalence classes, starting from the trivial partition and refining the previous partition with every step, until these partitions cannot be refined any further. We show that the equivalence classes produced in round n contain exactly the types that induce the same n -th order belief. In particular, the equivalence classes produced at the end contain precisely those types that induce the same (infinite) belief hierarchy.

Procedure 2 (Type partitioning procedure (finite type structures)). Consider a multi-agent uncertainty space $\mathcal{X} = (X_i, \Sigma_i)_{i \in I}$ and a finite type structure $\mathcal{T} = (T_i, \Sigma_i^T, b_i)_{i \in I}$ for \mathcal{X} .

Initial step: For every agent i , let \mathcal{P}_i^0 be the trivial partition of his or her set of types T_i .

⁷ This is with some abuse of notation, since b_i^* is defined on $X_i \times T_{-i}^*$, while b_i^1 and b_i^2 are defined on $X_i \times T_{-i}^1$ and $X_i \times T_{-i}^2$, respectively. By defining the σ -algebra $\Sigma_i^{T^*}$ on T_i^* as above, the extension of b_i^1 and b_i^2 to the larger domain is well defined.

Inductive step: Suppose that $n \geq 1$ and that the partitions \mathcal{P}_i^{n-1} have been defined for every agent i . Then, for every agent i , and every $t_i \in T_i$,

$$\mathcal{P}_i^n(t_i) = \{t'_i \in T_i \mid b_i(t'_i)(E_i \times P_{-i}^{n-1}) = b_i(t_i)(E_i \times P_{-i}^{n-1}) \quad (2)$$

for all $E_i \in \Sigma_i$, and all $P_{-i}^{n-1} \in \mathcal{P}_{-i}^{n-1}\}$.

The procedure terminates at round n whenever $\mathcal{P}_i^n = \mathcal{P}_i^{n-1}$ for every agent i .

In this procedure, \mathcal{P}_{-i}^{n-1} is the partition of the set T_{-i} induced by the partitions \mathcal{P}_j^{n-1} on T_j . Again, it follows from a simple inductive argument that \mathcal{P}_i^n is a refinement of \mathcal{P}_i^{n-1} for every player i and every n . Note that if the total number of types, viz., $\sum_{i \in I} |T_i|$, equals N , then the procedure terminates in at most $N - |I|$ steps. We now illustrate the procedure by means of an example.

Example 1. Consider the first type structure $\mathcal{T} = (T_1, T_2, \Sigma_1, \Sigma_2, b_1, b_2)$ from Table 1.

Initial step: Let \mathcal{P}_1^0 be the trivial partition of the set of types T_1 , and let \mathcal{P}_2^0 be the trivial partition of the set of types T_2 . That is,

$$\mathcal{P}_1^0 = \{\{t_1, t'_1, t''_1\}\} \text{ and } \mathcal{P}_2^0 = \{\{t_2, t'_2, t''_2\}\}.$$

Round 1: By Equation (2),

$$\begin{aligned} \mathcal{P}_1^1(t_1) &= \{\tau_1 \in T_1 \mid \\ b_1(\tau_1)(\{c\} \times T_2) &= b_1(t_1)(\{c\} \times T_2) = \frac{1}{2}, \\ b_1(\tau_1)(\{d\} \times T_2) &= b_1(t_1)(\{d\} \times T_2) = \frac{1}{2}\} \\ &= \{t_1, t'_1, t''_1\}, \end{aligned}$$

which implies that:

$$\mathcal{P}_1^1 = \mathcal{P}_1^0 = \{\{t_1, t'_1, t''_1\}\}.$$

At the same time,

$$\begin{aligned} \mathcal{P}_2^1(t_2) &= \{\tau_2 \in T_2 \mid \\ b_2(\tau_2)(\{e\} \times T_1) &= b_2(t_2)(\{e\} \times T_1) = \frac{3}{4}, \\ b_2(\tau_2)(\{f\} \times T_1) &= b_2(t_2)(\{f\} \times T_1) = \frac{1}{4}\} \\ &= \{t_2, t'_2\} \end{aligned}$$

which implies that $\mathcal{P}_2^1(t'_2) = \{t'_2\}$, and hence:

$$\mathcal{P}_2^1 = \{\{t_2, t'_2\}, \{t'_2\}\}.$$

Round 2: By Equation (2),

$$\begin{aligned} \mathcal{P}_1^2(t_1) &= \{\tau_1 \in T_1 \mid \\ b_1(\tau_1)(\{c\} \times \{t_2, t'_2\}) &= b_1(t_1)(\{c\} \times \{t_2, t'_2\}) = \frac{1}{2}, \\ b_1(\tau_1)(\{c\} \times \{t'_2\}) &= b_1(t_1)(\{c\} \times \{t'_2\}) = 0, \\ b_1(\tau_1)(\{d\} \times \{t_2, t'_2\}) &= b_1(t_1)(\{d\} \times \{t_2, t'_2\}) = 0, \\ b_1(\tau_1)(\{d\} \times \{t'_2\}) &= b_1(t_1)(\{d\} \times \{t'_2\}) = \frac{1}{2}\} \\ &= \{t_1, t'_1\}, \end{aligned}$$

which implies that $\mathcal{P}_1^2(t_1'') = \{t_1''\}$, and hence:

$$\mathcal{P}_1^2 = \{\{t_1, t_1'\}, \{t_1''\}\}.$$

Since $\mathcal{P}_1^1 = \mathcal{P}_1^0$, we may immediately conclude that:

$$\mathcal{P}_2^2 = \mathcal{P}_2^1 = \{\{t_2, t_2''\}, \{t_2'\}\}.$$

Round 3: As $\mathcal{P}_2^2 = \mathcal{P}_2^1$, we may immediately conclude that:

$$\mathcal{P}_1^3 = \mathcal{P}_1^2 = \{\{t_1, t_1'\}, \{t_1''\}\}.$$

By Equation (2),

$$\begin{aligned} \mathcal{P}_2^3(t_2) &= \{\tau_2 \in T_2 \mid \\ b_2(\tau_2)(\{e\} \times \{t_1, t_1'\}) &= b_2(t_2)(\{e\} \times \{t_1, t_1'\}) = \frac{3}{4}, \\ b_2(\tau_2)(\{e\} \times \{t_1''\}) &= b_2(t_2)(\{e\} \times \{t_1''\}) = 0, \\ b_2(\tau_2)(\{f\} \times \{t_1, t_1'\}) &= b_2(t_2)(\{f\} \times \{t_1, t_1'\}) = 0, \\ b_2(\tau_2)(\{f\} \times \{t_1''\}) &= b_2(t_2)(\{f\} \times \{t_1''\}) = \frac{1}{4} \} \\ &= \{t_2, t_2''\}, \end{aligned}$$

which implies that $\mathcal{P}_2^3(t_2') = \{t_2'\}$, and hence,

$$\mathcal{P}_2^3 = \{\{t_2, t_2''\}, \{t_2'\}\} = \mathcal{P}_2^2.$$

As $\mathcal{P}_1^3 = \mathcal{P}_1^2$ and $\mathcal{P}_2^3 = \mathcal{P}_2^2$, the procedure terminates at Round 3. The final partitions of the types are thus given by:

$$\mathcal{P}_1^\infty = \{\{t_1, t_1'\}, \{t_1''\}\} \text{ and } \mathcal{P}_2^\infty = \{\{t_2, t_2''\}, \{t_2'\}\}.$$

The reader may check that all types within the same equivalence class indeed induce the same belief hierarchy. That is, t_1 induces the same belief hierarchy as t_1' , and t_2 induces the same belief hierarchy as t_2'' . Moreover, t_1 and t_1'' induce different belief hierarchies, and so do t_2 and t_2' .

Our characterization result for the case of finite type structures states that the type partitioning procedure characterizes precisely those groups of types that induce the same belief hierarchy. We actually prove a little more: we show that the partitions generated in round n of the procedure characterize exactly those types that yield the same n -th order belief.

Proposition 5 (Characterization result (finite type structures)). *Consider a finite type structure $\mathcal{T} = (T_i, \Sigma_i, b_i)_{i \in I}$, where Σ_i is the discrete σ -algebra on T_i for every player i . For every agent i , every $n \geq 1$ and every two types $t_i, t_i' \in T_i$, we have that*

(a) $h_i^{\mathcal{T}, n}(t_i) = h_i^{\mathcal{T}, n}(t_i')$, if and only if, $t_i' \in \mathcal{P}_i^n(t_i)$;

(b) $h_i^{\mathcal{T}}(t_i) = h_i^{\mathcal{T}}(t_i')$, if and only if, $t_i' \in \mathcal{P}_i^\infty(t_i)$.

The proof follows directly from Proposition 3 and is therefore omitted. As before, this result can be used to verify whether two types from different type structures generate the same belief hierarchies, by first merging the two type structures and then running the type partitioning procedure on this “large” type structure.

Acknowledgments: This paper is a substantially revised version of an earlier paper with the same title by Andrés Perea. We would like to thank Pierpaolo Battigalli, Eddie Dekel, Amanda Friedenberg, Christian Nauerz, Miklós Pintér, Elias Tsakas and two anonymous reviewers for very useful comments.

Author Contributions: Both authors have contributed equally to this paper.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Proofs

Appendix A.1. Proof of Theorem 1

By definition, \mathcal{T}' contains \mathcal{T} if and only if there is a hierarchy morphism from \mathcal{T} to \mathcal{T}' . Therefore, it suffices to show that every generalized type morphism is a hierarchy morphism and vice versa.

Appendix A.1.1. Every Hierarchy Morphism Is a Generalized Type Morphism

To show that every hierarchy morphism is a generalized type morphism, we need to show two things. First, we need to show that any hierarchy morphism is measurable with respect to the appropriate σ -algebra. Second, we need to show that beliefs are preserved for the relevant events.

Let us start with the measurability condition. Suppose φ is a hierarchy morphism. Let $i \in I$ and $E \in \mathcal{F}_i^{\mathcal{T}'}$. We need to show that:

$$\{t_i \in T_i \mid \varphi_i(t_i) \in E\} \in \Sigma_i^{\mathcal{T}}.$$

Recall that $\mathcal{F}_i^{\mathcal{T}'} = \sigma(h_i^{\mathcal{T}'})$ (Lemma 1). Therefore, there is a measurable subset B of the set H_i of belief hierarchies, such that:

$$E = \{t'_i \in T'_i \mid h_i^{\mathcal{T}'}(t'_i) \in B\}.$$

Hence,

$$\begin{aligned} \{t_i \in T_i \mid \varphi_i(t_i) \in E\} &= \{t_i \in T_i \mid h_i^{\mathcal{T}'}(\varphi_i(t_i)) \in B\} \\ &= \{t_i \in T_i \mid h_i^{\mathcal{T}}(t_i) \in B\}, \end{aligned}$$

where the second equality follows from the assumption that φ is a hierarchy morphism. By Lemma 1, we have:

$$\{t_i \in T_i \mid h_i^{\mathcal{T}}(t_i) \in B\} \in \mathcal{F}_i^{\mathcal{T}}.$$

Since $\Sigma_i^{\mathcal{T}} \supseteq \mathcal{F}_i^{\mathcal{T}}$ (Definition 4 and Lemma 1), the result follows.

We next ask whether hierarchy morphisms preserve beliefs for the relevant events. Again, let φ be a hierarchy morphism. Let $i \in I$, $t_i \in T_i$ and $E' \in \Sigma_i \otimes \mathcal{F}_{-i}^{\mathcal{T}'}$. We need to show that:

$$b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1}(E') = b'_i(\varphi_i(t_i))(E'),$$

where Id_{X_i} is the identity function on X_i and where we have used the notation (f_1, \dots, f_m) for the induced function that maps (x_1, \dots, x_m) into $(f_1(x_1), \dots, f_m(x_m))$, so that $b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1}$ is the image measure induced by $(\text{Id}_{X_i}, \varphi_{-i})$. By a similar argument as before, there is a measurable subset B' of the set $X_i \times H_{-i}$, such that:

$$E' = \{(x_i, t'_{-i}) \in X_i \times T'_{-i} \mid (x_i, h_{-i}^{\mathcal{T}'}(t'_{-i})) \in B'\}.$$

If E' is an element of $\Sigma_i \otimes \bigotimes_{j \neq i} \{T'_j, \emptyset\}$, then the result follows directly from the definitions. Therefore, suppose $E' \notin \Sigma_i \otimes \bigotimes_{j \neq i} \{T'_j, \emptyset\}$. Then, for every $n \geq 1$, define:

$$\begin{aligned} B^n := \{(x_i, \mu_{-i}^1, \dots, \mu_{-i}^n) \in X_i \times H_{-i}^n \mid (x_i, \mu_{-i}^1, \dots, \mu_{-i}^n, \mu_{-i}^{n+1}, \dots) \in B' \\ \text{for some } (\mu_{-i}^{n+1}, \mu_{-i}^{n+2}, \dots)\} \end{aligned}$$

and:

$$E^n := \{(x_i, t'_i) \in X_i \times T'_i \mid (x_i, h_{-i}^{T',n}(t'_i)) \in B^n\}.$$

Then, $E^n \supseteq E'$ and $E^n \downarrow E'$. Furthermore, we have $E^n \in \Sigma_i \otimes \otimes_{j \neq i} \sigma(h_j^{T',n})$, and thus, $E^n \in \Sigma_i \otimes \mathcal{F}_{-i}^{T'}$ (Lemma 1). For every n ,

$$\begin{aligned} b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1}(E^{n-1}) &= b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1} \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1}(B^{n-1}) \\ &= b_i(t_i) \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1} \circ \varphi_{-i})^{-1}(B^{n-1}) \\ &= \mu_i^{T',n}(t_i)(B^{n-1}) \\ &= b'_i(\varphi(t_i)) \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1}(B^{n-1}) \\ &= b'_i(\varphi(t_i))(E^{n-1}), \end{aligned}$$

where the penultimate equality uses the definition of a hierarchy morphism. By the continuity of the probability measures $b_i(t_i)$ and $b'_i(\varphi_i(t_i))$ (e.g., [28], Thm. 10.8), we have $b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1}(E') = b'_i(\varphi_i(t_i))(E')$, and the result follows.

Appendix A.1.2. Every Generalized Type Morphism Is a Hierarchy Morphism

For the other direction, that is to show that every generalized type morphism is a hierarchy morphism, suppose that φ is a generalized type morphism from $\mathcal{T} = (T_i, \Sigma_i^{\mathcal{T}}, b_i)_{i \in I}$ to $\mathcal{T}' = (T'_i, \Sigma_i^{T'}, b'_i)_{i \in I}$. We can use an inductive argument to show that it is a hierarchy morphism. Let $i \in I$ and $t_i \in T_i$. Then, for all $E \in \Sigma_i$,

$$\begin{aligned} \mu_i^{T',1}(\varphi_i(t_i))(E) &= b'_i(\varphi_i(t_i))(E \times T'_{-i}) \\ &= b_i(t_i)(E \times T_{-i}) \\ &= \mu_i^{T,1}(t_i), \end{aligned}$$

where the first and the last equality use the definition of a first-order belief induced by a type and the second uses the definition of a generalized type morphism. Therefore, $\mu_i^{T',1}(t_i) = \mu_i^{T',1}(\varphi_i(t_i))$, and thus, $h_i^{T',1}(t_i) = h_i^{T',1}(\varphi_i(t_i))$ for each player i and every type $t_i \in T_i$.

For $n > 1$, suppose that for each player i and every type $t_i \in T_i$, we have $h_i^{T',n-1}(t_i) = h_i^{T',n-1}(\varphi_i(t_i))$. We will use the notation (f_1, \dots, f_m) for the induced function that maps (x_1, \dots, x_m) into $(f_1(x_1), \dots, f_m(x_m))$, so that $\mu \circ (f_1, \dots, f_m)^{-1}$ is the image measure induced by a probability measure μ and (f_1, \dots, f_m) .

Let E be a measurable subset of $X_i \times H_{-i}^n$. By Lemma 1, we have $\mathcal{F}_j^{T'} = \sigma(h_j^{T'})$; and clearly, $\sigma(h_j^{T'}) \supseteq \sigma(h_j^{T',n-1})$. Therefore, if we write Id_{X_i} for the identity function on X_i , we have $(\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1}(E) \in \Sigma_i \otimes \mathcal{F}_{-i}^{T'}$. Then, for every player i and type $t_i \in T_i$,

$$\begin{aligned} \mu_i^{T',n}(\varphi_i(t_i))(E) &= b'_i(\varphi_i(t_i)) \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1}(E) \\ &= b_i(t_i) \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1} \circ (\text{Id}_{X_i}, \varphi_{-i})^{-1}(E) \\ &= b_i(t_i) \circ (\text{Id}_{X_i}, \varphi_{-i} \circ h_{-i}^{T',n-1})^{-1}(E) \\ &= b_i(t_i) \circ (\text{Id}_{X_i}, h_{-i}^{T',n-1})^{-1}(E) \\ &= \mu_i^{T',n}(t_i)(E), \end{aligned}$$

where the first equality uses the definition of an n -th-order belief, the second uses the definition of a generalized type morphism, the third uses the definition of the composition operator, the fourth uses

the induction hypothesis and the fifth uses the definition of an n -th-order belief again. Conclude that $\mu_i^{\mathcal{T},n}(t_i) = \mu_i^{\mathcal{T},n}(\varphi_i(t_i))$ and thus $h_i^{\mathcal{T},n}(t_i) = h_i^{\mathcal{T},n}(\varphi_i(t_i))$ for each player i and every type $t_i \in T_i$.

Therefore, for each player $i \in I$ and each type $t_i \in T_i$, we have $h_i^{\mathcal{T}}(t_i) = h_i^{\mathcal{T}}(\varphi_i(t_i))$, which shows that φ is a hierarchy morphism. \square

Appendix A.2. Proof of Proposition 2

Let $i \in I$. It will be convenient to define $h_i^{\mathcal{T},0}$ to be the trivial function from T_i into some singleton $\{v_i\}$. Therefore, the σ -algebra $\sigma(h_i^{\mathcal{T},0})$ generated by $h_i^{\mathcal{T},0}$ is just the trivial σ -algebra $\mathcal{S}_i^0 = \{T_i, \emptyset\}$. Next, consider $n = 1$. Fix player $i \in I$. By definition, $\sigma(h_i^{\mathcal{T},1})$ is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid h_i^{\mathcal{T},1}(t_i) \in E\} : E \subseteq H_i^1 \text{ measurable.}$$

It suffices to restrict attention to the generating sets E of the σ -algebra on $H_i^1 = \Delta(X_i)$ (e.g., [28]). Therefore, $\sigma(h_i^{\mathcal{T},1})$ is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid h_i^{\mathcal{T},1}(t_i) \in E\}$$

where E is of the form $\{\mu \in \Delta(X_i) \mid \mu(F) \geq p\}$ for $F \in \Sigma_i$ and $p \in [0, 1]$. Using that for each type t_i , $h_i^{\mathcal{T},1}(t_i)$ is the marginal on X_i of $b_i(t_i)$, we have that $\sigma(h_i^{\mathcal{T},1})$ is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid b_i(t_i)(E) \geq p\} : E \in \Sigma_i \otimes \mathcal{S}_{-i}^0, p \in [0, 1].$$

That is, $\sigma(h_i^{\mathcal{T},1}) = \mathcal{S}_i^1$. In particular, $h_i^{\mathcal{T},1}$ is measurable with respect to $\mathcal{S}_i^{\mathcal{T},1}$.

For $n > 1$, suppose, inductively, that for each player $i \in I$, $\sigma(h_i^{\mathcal{T},n-1}) = \mathcal{S}_i^{\mathcal{T},n-1}$, so that $h_i^{\mathcal{T},n-1}$ is measurable with respect to $\mathcal{S}_i^{\mathcal{T},n-1}$. Fix $i \in I$. By definition, $\sigma(h_i^{\mathcal{T},n})$ is the coarsest σ -algebra that contains the sets in $\sigma(h_i^{\mathcal{T},n-1})$ and the sets:

$$\{t_i \in T_i \mid \mu_i^{\mathcal{T},n}(t_i) \in E\} : E \subseteq \Delta(X_i \times H_{-i}^{n-1}) \text{ measurable.}$$

Again, it suffices to consider the generating sets of the σ -algebra on $\Delta(X_i \times H_{-i}^{n-1})$. Hence, $\sigma(h_i^{\mathcal{T},n})$ is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i \mid \mu_i^{\mathcal{T},n}(t_i)(F) \geq p\} : F \subseteq X_i \times H_{-i}^{n-1} \text{ measurable and } p \in [0, 1].$$

(Note that this includes the generating sets of $\sigma(h_i^{\mathcal{T},n-1})$, given that the n -th-order belief induced by a type is consistent with its $(n - 1)$ -th-order belief.) Using the definition of $\mu_i^{\mathcal{T},n}$ and the induction assumption that $\mathcal{S}_{-i}^{\mathcal{T},n-1} = \sigma(h_{-i}^{\mathcal{T},n-1})$, we see that $\sigma(h_i^{\mathcal{T},n})$ is the coarsest σ -algebra on T_i that contains the sets:

$$\{t_i \in T_i \mid b_i(t_i)(F) \geq p\} : F \in \Sigma_i \otimes \mathcal{S}_{-i}^{\mathcal{T},n-1}, p \in [0, 1].$$

That is, $\sigma(h_i^{\mathcal{T},n}) = \mathcal{S}_i^{\mathcal{T},n}$, and $h_i^{\mathcal{T},n}$ is measurable with respect to $\mathcal{S}_i^{\mathcal{T},n}$.

Therefore, for each player i and $n \geq 1$, $\sigma(h_i^{\mathcal{T},n}) = \mathcal{S}_i^{\mathcal{T},n}$. It follows immediately that $\sigma(h_i^{\mathcal{T}})$, as the σ -algebra on T_i generated by the “cylinders” $\sigma(h_i^{\mathcal{T},n})$, is equal to $\mathcal{S}_i^{\mathcal{T},\infty}$. \square

Appendix A.3. Proof of Lemma 2

Let $i \in I$. Recall that \mathcal{X}_i is countably generated, that is there is a countable subset \mathcal{D}_i^0 of Σ_i , such that \mathcal{D}_i^0 generates Σ_i (i.e., Σ_i is the coarsest σ -algebra that contains \mathcal{D}_i^0). Throughout this proof, we write $\sigma(D)$ for the σ -algebra on a set Y generated by a collection D of subsets of Y .

The following result says that a countable collection of subsets of a set Y generates a countable algebra on Y . For a collection D of subsets of a set Y , denote the algebra generated by D by $A(D)$. Therefore, $A(D)$ is the coarsest algebra on Y that contains D .

Lemma 3. Let D be a countable collection of subsets of a set Y . Then, the algebra $A(D)$ generated by D is countable.

Proof. We can construct the algebra generated by D . Denote the elements of D by $D_\lambda, \lambda \in \Lambda$, where Λ is a countable index set. Define:

$$A(D) = \left\{ \bigcup_{m \in F} \bigcap_{\ell \in L_m} D_\ell, F \text{ a finite subset of } \mathbb{N}, L_m \text{ a finite subset of } \Lambda, \right. \\ \left. D_\ell = A_k \text{ or } D_\ell = A_k^c \text{ for some } k \right\}, \tag{A1}$$

where E^c is the complement of a set E . That is, $A(D)$ is the collection of finite unions of finite intersection of elements of D and their complements. We check that $A(D)$ is an algebra. Clearly, $A(D)$ is nonempty (it contains D) and $\emptyset \in A(D)$. We next show that $A(D)$ is closed under finite intersections. Let:

$$A_1 := \bigcup_{m_1 \in F_1} \bigcap_{\ell \in L_{m_1}^1} D_\ell, \quad A_2 := \bigcup_{m_2 \in F_2} \bigcap_{\ell \in L_{m_2}^2} D_\ell,$$

be elements of $A(D)$. Then,

$$A_1 \cap A_2 = \bigcup_{(m_1, m_2) \in F_1 \times F_2} \bigcap_{\ell_1 \in L_{m_1}^1} \bigcap_{\ell_2 \in L_{m_2}^2} D_{\ell_1} \cap D_{\ell_2}.$$

Clearly, $F_1 \times F_2$ is finite and so are the sets $L_{m_1}^1$ and $L_{m_2}^2$. We can thus rewrite $A_1 \cap A_2$ so that it is of the form as the elements in (A1). We can likewise show that $A(D)$ is closed under complements: let $A := \bigcup_{m \in F} \bigcap_{\ell \in L_m} D_\ell \in A(D)$, so that $A^c = \bigcap_{m \in F} \bigcup_{\ell \in L_m} D_\ell^c$; then, since $\bigcup_{\ell \in L_m} D_\ell^c \in A(D)$ for every m , we have $A^c \in A(D)$. Therefore, $A(D)$ is an algebra that contains D , and it is in fact the coarsest such one (by construction, any proper subset of $A(D)$ does not contain all finite intersections of the sets in D and their complements). As D is countable, so is the collection of the elements in D and their complements; the collections of the finite intersections of such sets are also countable. Hence, $A(D)$ is countable. \square

Note that for any $p \in [0, 1]$, the set $\{t_i \in T_i : b_i(t_i)(E) \geq p\}$ can be written as the countable intersection of sets $\{t_i \in T_i : b_i(t_i)(E) \geq p_\ell\}$ for some rational $p_\ell, \ell = 1, 2, \dots$. Therefore, by Proposition 2, the σ -algebra $\sigma(h_i^{T, n})$, $n = 1, 2, \dots$, on the type set $T_i, i \in I$, is the coarsest σ -algebra that contains the sets:

$$\{t_i \in T_i : b_i(t_i)(E) \geq p\} : \quad E \in \Sigma_i \otimes \bigotimes_{j \neq i} \sigma(h_j^{T, n-1}), p \in \mathbb{Q}$$

We are now ready to prove Lemma 2. Fix $i \in I$. By Lemma 3, the set \mathcal{D}_i^0 generates a countable algebra $A(\mathcal{D}_i^0)$ on X_i . Then, by Proposition 2 and by Lemma 4.5 of Heifetz and Samet [14], we have that the σ -algebra $\sigma(h_i^{T, 1})$ is generated by the sets:

$$\{t_i \in T_i : b_i(t_i)(E) \geq p\} : \quad E \in \mathcal{D}_i^0 \otimes \bigotimes_{j \neq i} \sigma(h_j^{T, 0}), p \in \mathbb{Q}.$$

Denote this collection of these sets by \mathcal{D}_i^1 , so that $\sigma(h_i^{T, 1}) = \sigma(\mathcal{D}_i^1)$; clearly, \mathcal{D}_i^1 is countable and $A(\mathcal{D}_i^1) \subseteq \sigma(h_i^{T, 1})$ (so that $\sigma(h_i^{T, 1}) = \sigma(A(\mathcal{D}_i^1))$).

For $m > 1$, suppose that for every $i \in I$, the σ -algebra $\sigma(h_i^{T, m-1})$ on T_i is generated by a countable collection \mathcal{D}_i^{m-1} of subsets of T_i , such that $A(\mathcal{D}_i^{m-1}) \subseteq \sigma(h_i^{T, m-1})$. Fix $i \in I$. By Proposition 2 and Lemma 4.5 of Heifetz and Samet [14], the σ -algebra $\sigma(h_i^{T, m})$ is generated by the sets:

$$\{t_i \in T_i : b_i(t_i)(E) \geq p\} : \quad E \in \mathcal{D}_i^0 \otimes \bigotimes_{j \neq i} A(\mathcal{D}_j^{m-1}), p \in \mathbb{Q}.$$

Denote this collection of these sets by \mathcal{D}_i^m ; as before, \mathcal{D}_i^m is clearly countable and $A(\mathcal{D}_i^m) \subseteq \sigma(h_i^{\mathcal{T},m})$. Again, we have $\sigma(h_i^{\mathcal{T},m}) = \sigma(\mathcal{D}_i^m) = \sigma(A(\mathcal{D}_i^m))$.

Therefore, we have shown that for every $i \in I$ and $m = 1, 2, \dots$, the σ -algebra $\sigma(h_i^{\mathcal{T},m})$ is generated by a countable collection \mathcal{D}_i^m of subsets of T_i . The σ -algebra $\sigma(h_i^{\mathcal{T}})$ is generated by the algebra $\bigcup_m \sigma(h_i^{\mathcal{T},m}) = \bigcup_m \sigma(\mathcal{D}_i^m)$ or, equivalently, by the union $\bigcup_m \mathcal{D}_i^m$ (Proposition 2). Since the latter set, as the countable union of countable sets, is countable, the σ -algebra $\sigma(h_i^{\mathcal{T}})$ is countably generated.

It now follows from Theorem V.2.1 of Parthasarathy [29] that for each player i , the σ -algebras $\sigma(h_i^{\mathcal{T},m})$, $m = 1, 2, \dots$ are atomic in the sense that for each $t_i \in T_i$, there is a unique atom $a_{t_i}^m$ in $\sigma(h_i^{\mathcal{T},m})$ containing t_i ; the analogous statement holds for $\sigma(h_i^{\mathcal{T}})$. \square

Appendix A.4. Proof of Proposition 3

Fix a player $i \in I$. By Proposition 2, we have $\mathcal{S}_i^{\mathcal{T},\infty} = \sigma(h_i^{\mathcal{T}})$ and $\mathcal{S}_i^{\mathcal{T},n} = \sigma(h_i^{\mathcal{T},n})$ for each $n \geq 1$. By Lemma 2, the σ -algebras $\sigma(h_i^{\mathcal{T},\infty})$ and $\sigma(h_i^{\mathcal{T},n})$ are atomic for every $n \geq 1$. Let $n \geq 1$. Let $t_i, t'_i \in T_i$. Since $\sigma(h_i^{\mathcal{T},n})$ is atomic, there exist a unique atom $a_{t_i}^n \in \sigma(h_i^{\mathcal{T},n})$, such that $t_i \in a_{t_i}^n$, and a unique atom $a_{t'_i}^n \in \sigma(h_i^{\mathcal{T},n})$, such that $t'_i \in a_{t'_i}^n$. Suppose $h_i^{\mathcal{T},n}(t_i) = h_i^{\mathcal{T},n}(t'_i)$. Then, for every generating set E of the σ -algebra $\sigma(h_i^{\mathcal{T},n})$, either $t_i, t'_i \in E$ or $t_i, t'_i \notin E$. Therefore, $a_{t_i}^n = a_{t'_i}^n$. Suppose $h_i^{\mathcal{T}}(t_i) \neq h_i^{\mathcal{T}}(t'_i)$. Then, there is a generating set E of $\sigma(h_i^{\mathcal{T},n})$ that separates t_i and t'_i , that is, $t_i \in E$, $t'_i \notin E$. Therefore, $a_{t_i}^n \neq a_{t'_i}^n$. The proof of the claim that there is a unique atom $a_{t_i}^\infty$ in $\sigma(h_i^{\mathcal{T}})$ that contains both t_i and t'_i if and only if $h_i^{\mathcal{T}}(t_i) = h_i^{\mathcal{T}}(t'_i)$ is analogous and therefore omitted. \square

References

- Carlsson, H.; van Damme, E. Global Games and Equilibrium Selection. *Econometrica* **1993**, *61*, 989–1018.
- Feinberg, Y.; Skrzypacz, A. Uncertainty about uncertainty and delay in bargaining. *Econometrica* **2005**, *73*, 69–91.
- Friedenberg, A. *Bargaining under Strategic Uncertainty*; Working Paper; Arizona State University: Tempe, AZ, USA, 2014.
- Geanakoplos, J.D.; Polemarchakis, H.M. We can't disagree forever. *J. Econ. Theory* **1982**, *28*, 192–200.
- Neeman, Z. The relevance of private information in mechanism design. *J. Econ. Theory* **2004**, *117*, 55–77.
- Brandenburger, A.; Dekel, E. Rationalizability and Correlated Equilibria. *Econometrica* **1987**, *55*, 1391–1402.
- Tan, T.; Werlang, S. The Bayesian foundations of solution concepts in games. *J. Econ. Theory* **1988**, *45*, 370–391.
- Aumann, R.; Brandenburger, A. Epistemic conditions for Nash equilibrium. *Econometrica* **1995**, *63*, 1161–1180.
- Perea, A. A one-person doxastic characterization of Nash strategies. *Synthese* **2007**, *158*, 251–271.
- Battigalli, P.; Siniscalchi, M. Strong belief and forward-induction reasoning. *J. Econ. Theory* **2002**, *106*, 356–391.
- Harsanyi, J.C. Games on incomplete information played by Bayesian players. Part I. *Manag. Sci.* **1967**, *14*, 159–182.
- Friedenberg, A.; Meier, M. On the relationship between hierarchy and type morphisms. *Econ. Theory* **2011**, *46*, 377–399.
- Mertens, J.F.; Zamir, S. Formulation of Bayesian analysis for games with incomplete information. *Int. J. Game Theory* **1985**, *14*, 1–29.
- Heifetz, A.; Samet, D. Topology-Free Typology of Beliefs. *J. Econ. Theory* **1998**, *82*, 324–341.
- Dekel, E.; Fudenberg, D.; Morris, S. Interim Correlated Rationalizability. *Theor. Econ.* **2007**, *2*, 15–40.
- Ely, J.; Peski, M. Hierarchies of beliefs and interim rationalizability. *Theor. Econ.* **2006**, *1*, 19–65.
- Liu, Q. On redundant types and Bayesian formulation of incomplete information. *J. Econ. Theory* **2009**, *144*, 2115–2145.
- Friedenberg, A.; Meier, M. The context of the game. *Econ. Theory* **2016**, in press.
- Yildiz, M. Invariance to Representation of Information. *Games Econ. Behav.* **2015**, *94*, 142–156.
- Brandenburger, A.; Keisler, H.J. An Impossibility Theorem on Beliefs in Games. *Stud. Log.* **2006**, *84*, 211–240.
- Friedenberg, A. When Do Type Structures Contain All Hierarchies of Beliefs? *Games Econ. Behav.* **2010**, *68*, 108–129.

22. Kets, W. *Bounded Reasoning and Higher-Order Uncertainty*; Working Paper; Northwestern University: Evanston, IL, USA, 2011.
23. Friedenberg, A.; Keisler, H.J. *Iterated Dominance Revisited*; Working Paper; Arizona State University: Tempe, AZ, USA; University of Wisconsin-Madison: Madison, WI, USA, 2011.
24. Heifetz, A.; Samet, D. Knowledge Spaces with Arbitrarily High Rank. *Games Econ. Behav.* **1998**, *22*, 260–273.
25. Battigalli, P.; Siniscalchi, M. Hierarchies of conditional beliefs and interactive epistemology in dynamic games. *J. Econ. Theory* **1999**, *88*, 188–230.
26. Blume, L.; Brandenburger, A.; Dekel, E. Lexicographic Probabilities and Choice Under Uncertainty. *Econometrica* **1991**, *59*, 61–79.
27. Heifetz, A.; Kets, W. *Robust Multiplicity with a Grain of Naiveté*; Working Paper; Northwestern University: Evanston, IL, USA, 2013.
28. Aliprantis, C.D.; Border, K.C. *Infinite Dimensional Analysis: A Hitchhiker's Guide*, 3rd ed.; Springer: Berlin, Germany, 2005.
29. Parthasarathy, K. *Probability Measures on Metric Spaces*; AMS Chelsea Publishing: Providence, RI, USA, 2005.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).