



Reasoning about your own future mistakes [☆]Martin Meier ^a, Andrés Perea ^{b, , }*^a Free University of Bozen-Bolzano, Faculty of Economics and Management, Italy^b Maastricht University, Department of Quantitative Economics and EpiCenter, the Netherlands

ARTICLE INFO

JEL classification:
C72

Keywords:

Dynamic games
Perfect backwards rationalizability
Perfect quasi-perfect equilibrium
Perfect rationalizability
Procedural quasi-perfect rationalizability
Quasi-perfect rationalizability
Non-standard analysis
Cautious reasoning
Own mistakes

ABSTRACT

We propose a model of reasoning in dynamic games in which a player, at each information set, holds a conditional belief about his own future choices and the opponents' future choices. These conditional beliefs are assumed to be *cautious*, that is, the player never completely rules out any feasible future choice by himself or the opponents. We impose the following key conditions: (a) a player always believes that he will choose rationally in the future, (b) a player always believes that his opponents will choose rationally in the future, and (c) a player deems his own mistakes infinitely less likely than the opponents' mistakes. These conditions, together with iterating property (b), lead to the new concept of *perfect backwards rationalizability*. We show that perfect backwards rationalizable strategies exist in every finite dynamic game. We prove, moreover, that perfect backwards rationalizability constitutes a refinement of both *perfect rationalizability* (a rationalizability analogue to Selten's (1975) perfect equilibrium) and *procedural quasi-perfect rationalizability* (a rationalizability analogue to van Damme's (1984) quasi-perfect equilibrium) – two concepts that are introduced in this paper. As a consequence, our concept avoids both weakly dominated strategies in the normal form and strategies containing weakly dominated actions in the agent normal form. For one-shot games, the concept coincides with *permissibility* (Brandenburger (1992), Börgers (1994)).

1. Introduction

When reasoning in a dynamic game, it is natural to assume that your opponents – and even you yourself – may make mistakes with some small probability. We can formalize this idea by requiring that a player, at each of his information sets, assigns a – possibly infinitesimal – strictly positive probability to each of the opponents' choices and to each of his own future choices. To see how this may affect the strategy choice of a player, consider the game Γ^1 in the left-hand panel of Fig. 1, where player 1 is the only player.

If player 1 fears that, by mistake, he may choose d in the future, then his unique best choice is to go for strategy b .

However, without further restrictions, the assumptions above may still leave room for intuitively implausible behavior. Consider, for instance, the game Γ^2 in the right-hand panel of Fig. 1. In view of the assumptions above, player 1 is free to believe that his own future mistake d is more likely than player 2's mistake f , which allows player 1 to go for strategy b . At the same time, (a, c) seems the only plausible strategy for player 1, since believing that your own future mistakes are more likely than those of your opponents strikes us as rather counterintuitive.

[☆] We thank an associate editor and an anonymous referee for their helpful suggestions which helped us to substantially improve the paper. We are also grateful to some associate editors and anonymous referees for their valuable feedback on earlier versions of this paper. Finally, we would like to thank the participants at TARK 2019 for their useful comments.

* Corresponding author.

E-mail addresses: Martin.Meier@unibz.it (M. Meier), a.perea@maastrichtuniversity.nl (A. Perea).

<https://doi.org/10.1016/j.geb.2025.09.010>

Received 6 September 2024




Fig. 1. Belief about your own future mistakes.

To rule out such implausible behavior, we additionally require a player to deem his own future mistakes *infinitely less likely* than the mistakes by his opponents. The main goal of this paper is to develop a *rationalizability* concept for dynamic games, termed *perfect backwards rationalizability*¹, that satisfies this extra condition, together with the assumptions above.

More precisely, in perfect backwards rationalizability a player holds, at each of his information sets, a cautious belief about his opponents' strategies and a cautious belief about his own continuation strategy. Here, by *cautious* we mean that the player never rules out any feasible opponent's strategy, and never rules out any of his own continuation strategies, although he may deem some of these strategies infinitely more likely than other strategies. This is similar to the reasoning in *perfect equilibrium* (Selten (1975)) where a player always believes that his opponents, and he himself, will make mistakes with small probabilities.

We then impose the following key conditions on these beliefs: (a) a player always believes that he will himself choose rationally in the future, given his future beliefs about the opponents' strategies and his future beliefs about himself; (b) a player always believes that his opponents will choose rationally in the future; and (c) a player always deems his own future mistakes infinitely less likely than the mistakes by others. In (a) and (b), when we say that a player believes that he, or another player, chooses rationally, we mean that he only assigns infinitesimal probability to all suboptimal strategies. By imposing the properties (a) and (c), and iterating the property (b), we arrive at the concept of perfect backwards rationalizability.

With respect to condition (c), note that mistakes are not about ability but about unintended deviations from the planned choices. Since, by introspection, you know yourself better than others, it seems natural to deem your own mistakes less likely than those of your opponents. One could consider a weaker concept in which a player believes that his own mistakes are less likely, but not necessarily *infinitely* less likely, than those of the opponents. In this paper we adopt the stronger version.

Formally, we define the concept of perfect backwards rationalizability by the recursive elimination of strategies and beliefs in the game (see Section 3).² In Theorem 3.1 we show that this elimination procedure always yields a non-empty output in every finite dynamic game, and hence perfect backwards rationalizable strategies always exist. The existence proof, which can be found in Appendix B, is constructive.

Property (b) shows that, in terms of reasoning, perfect backwards rationalizability is very similar to *backwards rationalizability* (Perea (2014), Penta (2015)), in which players also always believe that their opponents will choose rationally in the future. In fact, we show in Section 4.2 that perfect backwards rationalizability always provides a refinement of backwards rationalizability.

By virtue of the properties (a), (b) and (c) above, especially property (c), the concept of perfect backwards rationalizability uniquely selects the “right” choice for player 1 in both of the games Γ^1 and Γ^2 above. Indeed, in game Γ^1 perfect backwards rationalizability uniquely selects strategy *b* for player 1, as player 1 does not rule out his own future mistake *d*. The selection here thus coincides with that of perfect equilibrium. In game Γ^2 , player 1's reasoning in line with perfect backwards rationalizability is as follows: Player 1 deems it possible that he himself will make the mistake *d* and that player 2 will make the mistake *f*, but by (c) he deems player 2's mistake infinitely more likely than his own mistake. Therefore, player 1 will go for strategy (*a, c*), which is also the prediction of quasi-perfect equilibrium (van Damme (1984)) and quasi-perfect rationalizability (Asheim and Perea (2005)).

In this paper we show that this is not a coincidence: In every dynamic game, perfect backwards rationalizability is a refinement of both *perfect rationalizability* (see Remark 4.1) and *procedural quasi-perfect rationalizability* (see Theorem 4.1) – two new concepts that are introduced in this paper. Here, perfect rationalizability may be viewed as a rationalizability analogue to perfect equilibrium, whereas procedural quasi-perfect rationalizability is a slight weakening of quasi-perfect rationalizability, and may be seen as a rationalizability analogue to quasi-perfect equilibrium. The main difference between the two versions of quasi-perfect rationalizability is that the former is defined by means of a recursive procedure, whereas the latter is characterized by epistemic conditions. But both concepts are based on the same principles: a player always believes in the opponents' future rationality, believes that the opponents make

¹ In a previous version of this paper, the concept was called *perfect quasi-perfect rationalizability*.

² A problem we would like to explore in the near future is characterizing perfect backwards rationalizability *epistemically*. That is, what are precisely the epistemic conditions on the players' beliefs that lead us to perfect backwards rationalizability? Our current formulation of this concept is by means of a non-epistemic recursive elimination procedure, although the conditions of “belief in your own future rationality” and “deeming your own mistakes least likely” already have some epistemic flavor.

Table 1
Cautious reasoning concepts discussed in this paper.

	Equilibrium	Rationalizability
Dynamic Games	Perfect Equilibrium (Selten (1975))	Perfect Rationalizability (*)
	Quasi-Perfect Equilibrium (van Damme (1984))	Quasi-Perfect Rationalizability (Asheim and Perea (2005)) Procedural Quasi-Perfect Rationalizability (*)
	Perfect Quasi-Perfect Equilibrium (Blume and Meier (2019))	Perfect Backwards Rationalizability (*)
Static Games	Perfect Equilibrium (Selten (1975))	Permissibility (Brandenburger (1992), Börgers (1994))

mistakes with infinitesimal probability, but believes that he himself will make no mistakes in the future. Typically, both concepts coincide behaviorally, but there are dynamic games where the former is behaviorally weaker than the latter.

The two results mentioned above thus show that, in every dynamic game, perfect backwards rationalizability inherits the desirable properties of both perfect rationalizability (that you take into account your own future mistakes) and procedural quasi-perfect rationalizability (that you focus primarily on the *opponents'* mistakes). As a consequence, perfect backwards rationalizability rules out strategies containing weakly dominated *actions* in the agent normal form, as well as weakly dominated *strategies* in the normal form. This follows from the fact that perfect rationalizability avoids strategies containing weakly dominated actions in the agent normal form, and procedural quasi-perfect rationalizability avoids weakly dominated strategies in the normal form. In a one-shot game, where the normal form and agent normal form coincide, it is shown that perfect backwards rationalizability is equivalent to the concept of *permissibility* (Brandenburger (1992), Börgers (1994)), which may be viewed as a rationalizability analogue to perfect equilibrium in one-shot games.

Remarkably, the result that perfect backwards rationalizability is a refinement of both perfect rationalizability and procedural quasi-perfect rationalizability does not have an equilibrium counterpart. Blume and Meier (2019) propose the concept of *perfect quasi-perfect equilibrium*, which may be viewed as an equilibrium analogue to perfect backwards rationalizability. They show that it is a refinement of quasi-perfect equilibrium, but not of perfect equilibrium. In fact, it cannot be a refinement of both as the sets of quasi-perfect equilibria and perfect equilibria may be disjoint, as shown in Blume and Meier (2019).

In dynamic games with perfect information, the differences between the various cautious reasoning concepts listed above only appear when there are relevant ties. For if we consider a dynamic game with perfect information without relevant ties, then all the above mentioned concepts (except for permissibility) would single out the unique backward induction strategy for each of the players.

Table 1 provides an overview of the various cautious reasoning concepts we discuss in this paper, and indicates when one concept can be viewed as an equilibrium, or rationalizability, analogue of another concept. The concepts with an asterisk are introduced in this paper.

The outline of this paper is as follows. In Section 2 we introduce the necessary notation for dynamic games and provide a brief overview of non-standard analysis, which will be sufficient for understanding the main body of this paper. In Section 3 we introduce the new concept of perfect backwards rationalizability and show its existence. In Section 4 we show how the concept relates to perfect rationalizability, procedural quasi-perfect rationalizability, and permissibility. In Section 5 we discuss why we opt for non-standard analysis instead of lexicographic beliefs for defining our concept, and how the concept relates to quasi-perfect rationalizability and perfect quasi-perfect equilibrium. Section 6 concludes with some final remarks. Appendix A gives a more extensive treatment of non-standard probabilities, which is needed for some of the proofs. Appendices B and C contain the proofs for Sections 3 and 4, respectively. Appendix D, finally, explores the relation between our notion of procedural quasi-perfect rationalizability and quasi-perfect rationalizability as defined in Asheim and Perea (2005).

2. Preliminaries

In this section we introduce the notation for dynamic games, and provide a short overview of non-standard analysis which is needed for our definition of *perfect backwards rationalizability* and other cautious reasoning concepts in this paper.

2.1. Dynamic games

In this paper we will focus on finite dynamic games with complete information that allow for simultaneous moves and imperfect information. The restriction to complete information is merely for the sake of simplicity. The definition below, and the concept presented in this paper, can easily be extended to games with incomplete information. Moreover, to keep our notation and definitions simple we exclude moves of nature in the definition that follows. However, our definition can easily be generalized to situations that involve moves of nature. Formally, a *finite dynamic game* is a tuple

$$G = (I, X, Z, (X_i)_{i \in I}, (C_i(x))_{i \in I, x \in X_i}, (H_i)_{i \in I}, (u_i)_{i \in I})$$

where

- (a) $I = \{1, 2, \dots, n\}$ is the finite set of *players*;

(b) X is the finite set of *histories*, consisting of *non-terminal* and *terminal* histories. At every non-terminal history, one or more players must make a choice, whereas at every terminal history the game ends. By \emptyset we denote the history that marks the beginning of the game;

(c) $Z \subseteq X$ is the set of *terminal* histories;

(d) $X_i \subseteq X$ is the set of *non-terminal* histories where player i must make a choice. For a given non-terminal history x , we denote by $I(x) := \{i \in I \mid x \in X_i\}$ the set of *active* players at x . We allow $I(x)$ to contain more than one player, that is, we allow for *simultaneous* moves. At the same time, we require $I(x)$ to be non-empty for every non-terminal history x ;

(e) $C_i(x)$ is the finite set of *choices* available to player i at a history $x \in X_i$;

(f) H_i is the collection of *information sets* for player i . Every information set $h \in H_i$ is a subset of histories in X_i such that $\cup_{h \in H_i} h = X_i$ and $h \cap h' = \emptyset$ for every two different $h, h' \in H_i$. That is, H_i is a partition of X_i . Moreover, we assume that $C_i(x) = C_i(y)$ whenever x, y belong to the same information set in H_i , and $C_i(x) \cap C_i(y) = \emptyset$ whenever x and y belong to different information sets in H_i . The interpretation of an information set $h \in H_i$ is that player i at h knows that a history in h has been realized. However, if h contains more than one history, player i does not know which of these histories has been realized. Hence, we allow for *imperfect information*;

(g) $u_i : Z \rightarrow \mathbb{R}$ is player i 's *utility function*, assigning to every terminal history $z \in Z$ some utility $u_i(z)$.

For practical purposes, we assume that all players are active at the beginning of the game \emptyset , that is, $I(\emptyset) = I$. If, in reality, player i does not choose at \emptyset , then we define $C_i(\emptyset)$ to be a singleton.

For every non-terminal history x and choice combination $(c_i)_{i \in I(x)}$ in $\times_{i \in I(x)} C_i(x)$, we denote by $x' = (x, (c_i)_{i \in I(x)})$ the (terminal or non-terminal) history that immediately follows this choice combination at x . In this case, we say that x' immediately follows x . We say that a history x follows a non-terminal history x' if there is a sequence of histories x^1, \dots, x^K such that $x^1 = x'$, $x^K = x$, and x^{k+1} immediately follows x^k for all $k \in \{1, \dots, K - 1\}$. A history x is said to *weakly follow* x' if either x follows x' or $x = x'$. In the obvious way, we can then also define what it means for x to (weakly) precede another history x' . Analogously, for two information sets h and h' , we say that h (weakly) follows h' if there is some $x \in h$ and $x' \in h'$ such that x (weakly) follows x' . We assume that the dynamic game has *non-overlapping information sets*, that is, for every two information sets h, h' it is never the case that h weakly follows h' and h' follows h .

In view of (f), we can write $C_i(h)$ to denote the (unique) set of choices that player i has available at information set $h \in H_i$. We assume *perfect recall*, that is, for every information set $h \in H_i$ and every two histories $x, y \in H_i$, the sequence of player i choices leading to x is the same as the sequence of player i choices leading to y . In particular, since different information sets in H_i prescribe disjoint sets of available choices, the sequence of player i information sets on the path to x and on the path to y must be the same. That is, player i always remembers, at each of his information sets $h \in H_i$, the choices he made in the past and the information he had in the past.

We view a strategy for player i as a *plan of action* (Rubinstein (1991)), assigning choices only to those histories $h \in H_i$ that are not precluded by previous choices. Formally, consider a collection of information sets $\hat{H}_i \subseteq H_i$, and a mapping $s_i : \hat{H}_i \rightarrow \cup_{h \in \hat{H}_i} C_i(h)$ assigning to every information set $h \in \hat{H}_i$ some available choice $s_i(h) \in C_i(h)$. We say that an information set $h \in H$ is *reachable* under s_i if at every information set $h' \in \hat{H}_i$ preceding h , the choice $s_i(h')$ is the unique choice that leads to h . The mapping $s_i : \hat{H}_i \rightarrow \cup_{h \in \hat{H}_i} C_i(h)$ is called a *strategy* if \hat{H}_i contains exactly those information sets in H_i that are reachable under s_i .³

By S_i we denote the set of strategies for player i . For every information set $h \in H$ and player i , we denote by $S_i(h)$ the set of strategies for player i under which h is reachable. Similarly, for a given strategy s_i we denote by $H_i(s_i)$ the collection of information sets in H_i that are reachable under s_i .

2.2. Non-standard numbers

The analysis of non-standard numbers was initiated by Robinson (1973). It has later been incorporated into the analysis of games by Hammond (1994) and Halpern (2010), who review non-standard probabilities and connect these to conditional and lexicographic probability systems.

Consider a number ε called an *infinitesimal*. Following Robinson (1973), Hammond (1994) and Halpern (2010), let $\mathbb{R}(\varepsilon)$ be the smallest field that includes all real numbers and the infinitesimal ε . That is, $\mathbb{R}(\varepsilon)$ contains all numbers a that can be written as

$$a = \frac{a_0 + a_1\varepsilon + a_2\varepsilon^2 + \dots + a_K\varepsilon^K}{b_0 + b_1\varepsilon + b_2\varepsilon^2 + \dots + b_K\varepsilon^K},$$

where $a_k, b_k \in \mathbb{R}$ for all $k \in \{0, \dots, K\}$, $b_k \neq 0$ for some $k \in \{0, \dots, K\}$, and where either $a_0 \neq 0$ or $b_0 \neq 0$. In other words, $\mathbb{R}(\varepsilon)$ contains all fractions of finite polynomials in ε . Numbers in $\mathbb{R}(\varepsilon)$ are called *non-standard*.

A non-standard number $a \in \mathbb{R}(\varepsilon)$ is *finite* if $b_0 \neq 0$. Every finite non-standard number $a \in \mathbb{R}(\varepsilon)$ can uniquely be written as

$$a = a_0 + a_1\varepsilon + a_2\varepsilon^2 + \dots,$$

³ Note that a strategy, by construction, specifies the planned choices in the absence of mistakes. At the same time, as we will see in Section 3, the concept of perfect backwards rationalizability requires a player to hold beliefs about his own future choices after possible mistakes by himself.

where $a_k \in \mathbb{R}$ for every $k \in \mathbb{N}_0$ (see Appendix A for a proof). Here, we use the notation $\mathbb{N} := \{1, 2, \dots\}$ and $\mathbb{N}_0 := \{0, 1, \dots\}$. By $st(a) := a_0$ we denote the *standard part* of a , which is the real number that is “closest” to a . By $v(a)$ we denote the smallest index $k \in \mathbb{N}_0$ for which $a_k \neq 0$, and call it the *valuation* of a . We say that $a = 0$ if $a_k = 0$ for all $k \in \mathbb{N}_0$, and we write $a > 0$ if $a_{v(a)} > 0$. For every two finite numbers $a, b \in \mathbb{R}(\varepsilon)$ we write $a > b$ if $a - b > 0$. For two finite numbers $a, b \in \mathbb{R}(\varepsilon)$, we say that a is of *infinitely smaller size* than b if $v(a) > v(b)$. We use the term “infinitely smaller size” rather than the more familiar “infinitely smaller” because we also apply it to negative numbers. For instance, ε is of infinitely smaller size than -1 , although ε is not smaller than -1 .

2.3. Non-standard probability distributions

Consider a finite set X . A *non-standard probability distribution* on X is a function $p : X \rightarrow \mathbb{R}(\varepsilon)$ such that $p(x) \geq 0$ for all $x \in X$ and $\sum_{x \in X} p(x) = 1$. We say that p is *cautious* on X if $p(x) > 0$ for all $x \in X$. We say that p *believes* an event $E \subseteq X$ if $\sum_{x \in E} p(x)$ has standard part 1. Consider, for instance, the set $X = \{x, y, z\}$ and the non-standard probability distribution p on X with $p(x) = 1 - \varepsilon - \varepsilon^2$, $p(y) = \varepsilon$ and $p(z) = \varepsilon^2$. Note that p is cautious, since every probability is strictly positive. Moreover, ε is of infinitely smaller size than $1 - \varepsilon - \varepsilon^2$, and ε^2 is of infinitely smaller size than ε . Hence, p can be interpreted as a cautious belief in which you deem event $\{x\}$ infinitely more likely than $\{y\}$, and event $\{y\}$ infinitely more likely than $\{z\}$, while deeming each of these three events possible. Note that p believes the event $\{x\}$.

For a subset $Y \subseteq X$ with $\sum_{x \in Y} p(x) > 0$, the *conditional probability distribution* on Y induced by p is the non-standard probability distribution p_Y on Y given by

$$p_Y(x) := \frac{p(x)}{\sum_{y \in Y} p(y)}$$

for every $x \in Y$.

A more extensive treatment of non-standard probabilities, which is needed for some of the proofs, can be found in Appendix A.

3. Perfect backwards rationalizability

The main ideas in *perfect backwards rationalizability* are that a player (a) is cautious about the opponents’ behavior and his own behavior, that is, he assigns a – possibly infinitesimal – strictly positive probability to every opponent’s strategy and every continuation strategy by himself, (b) always believes that his opponents and he himself will choose rationally in the future, and (c) believes that he may make mistakes himself in the future, but deems his own future mistakes much less likely – in fact, infinitely less likely – than the mistakes by his opponents.

Formally, for every player i , let B_i^{self} be the set of cautious non-standard probability distributions on the set S_i of i ’s own strategies. A member $b_i^{self} \in B_i^{self}$ will be interpreted as a *belief* that player i has about his own future choices in the game. Hence, every belief in B_i^{self} always deems each of his own future choices possible. The assumption that a player holds a belief about his own future choices seems natural, but is certainly not standard in game theory. Battigalli et al. (2013) and Battigalli and De Vito (2021) are some of the few papers that incorporate beliefs about own choices in a game-theoretic setting.

Moreover, let B_i^{opp} be the set of cautious non-standard probability distributions on the set S_{-i} of opponents’ strategy combinations, where $S_{-i} := \times_{j \neq i} S_j$. A member $b_i^{opp} \in B_i^{opp}$ represents a *belief* of player i about the opponents’ strategies. By definition, every belief in B_i^{opp} deems each of the opponents’ strategy combinations possible. By B_i we denote the set of belief pairs $b_i = (b_i^{self}, b_i^{opp})$ where $b_i^{self} \in B_i^{self}$ and $b_i^{opp} \in B_i^{opp}$.

Consider an information set $h \in H_i$ and a choice $c_i \in C_i(h)$ available at h . By $S_i(h, c_i)$ we denote the set of strategies $s_i \in S_i(h)$ with $s_i(h) = c_i$. By

$$S_{-i}(h) := \{s_{-i} \in S_{-i} \mid \text{there is some } s_i \in S_i \text{ such that } (s_i, s_{-i}) \text{ reaches a history in } h\}$$

we denote the set of opponents’ strategy combinations that are possible when h is reached. For a given belief pair $b_i = (b_i^{self}, b_i^{opp})$, let $b_i^{self}(h, c_i)$ be the induced conditional belief on $S_i(h, c_i)$, and let $b_i^{opp}(h)$ be the induced conditional belief on $S_{-i}(h)$. By

$$u_i(c_i, b_i, h) := \sum_{s_i \in S_i(h, c_i)} \sum_{s_{-i} \in S_{-i}(h)} b_i^{self}(h, c_i)(s_i) \cdot b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i}))$$

we denote the expected (non-standard) utility at information set h of making choice c_i under the belief pair b_i . Here, $z(s_i, s_{-i})$ is the outcome reached by the strategy combination (s_i, s_{-i}) .

A choice $c_i \in C_i(h)$ is *locally rational* for the belief pair $b_i = (b_i^{self}, b_i^{opp})$ at information set $h \in H_i$ if

$$u_i(c_i, b_i, h) \geq u_i(c'_i, b_i, h) \text{ for all } c'_i \in C_i(h).$$

For every information set h we define

$$S_i^{rat}(b_i, h) := \{s_i \in S_i(h) \mid s_i(h') \text{ locally rational for } b_i \text{ at every } h' \in H_i(s_i) \text{ following } h\}.$$

We say that the belief pair $b_i = (b_i^{self}, b_i^{opp})$ believes in his own future rationality if for every information set $h \in H_i$ and every choice $c_i \in C_i(h)$, the induced conditional belief $b_i^{self}(h, c_i)$ believes $S_i^{rat}(b_i, h)$. Note that at information set $h \in H_i$, player i need not believe that his choice c_i at h is optimal. Indeed, it may well be that c_i is suboptimal. The definition above only requires player i to believe at $h \in H_i$ that his own choices strictly following h are optimal. This condition is similar to the notion of *optimal planning* in Battigalli and De Vito (2021).

We say that b_i deems his own mistakes least likely⁴ if for every information set $h \in H_i$, every choice $c_i \in C_i(h)$, every strategy $s_i \in S_i(h, c_i)$ and every strategy combination $s_{-i} \in S_{-i}$ we have that

$$v[b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i)(s_i))] > v(b_i^{opp}(s_{-i})).$$

Recall from Section 2.2 that for a non-standard number $a = a_0 + a_1\varepsilon + a_2\varepsilon^2 + \dots$, the valuation $v(a)$ is the lowest index k with $a_k \neq 0$. That is, the infinitesimal mistake part $b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i)(s_i))$ in the belief about i 's own strategy is of infinitely smaller size than each of the belief probabilities $b_i^{opp}(s_{-i})$ about the opponent's strategies.

We now define *perfect backwards rationalizability* by recursively defining sets of strategies $S_i^k(h)$, for every information set h , and sets of belief pairs B_i^k , as follows. Remember that \emptyset is the information set that marks the beginning of the game.

Definition 3.1 (*Perfect backwards rationalizability*). (Initial step) Set $S_i^0(h) := S_i(h)$ and

$$B_i^0 := \{b_i \in B_i \mid b_i \text{ believes in his own future rationality and deems own mistakes least likely}\}$$

for all players i and all information sets h .⁵

(Inductive step) Let $k \in \mathbb{N}$, and suppose that $S_i^{k-1}(h)$ and B_i^{k-1} have been defined for all players i and all information sets h . Then define, for every player i and every information set h , the sets

$$S_i^k(h) := \{s_i \in S_i^{k-1}(h) \mid \text{there is some } b_i \in B_i^{k-1} \text{ such that } s_i(h') \text{ is locally rational for } b_i \text{ at every } h' \in H_i(s_i) \text{ weakly following } h\},$$

$$S_{-i}^k(h) := \{(s_j)_{j \neq i} \in S_{-i}(h) \mid s_j \in S_j^k(h) \text{ for all } j \neq i\},$$

and

$$B_i^k := \{b_i = (b_i^{self}, b_i^{opp}) \in B_i^{k-1} \mid b_i^{opp}(h) \text{ believes } S_{-i}^k(h) \text{ for all } h \in H_i\}.$$

A strategy $s_i \in S_i$ is called perfect backwards rationalizable if $s_i \in S_i^k(\emptyset)$ for all $k \in \mathbb{N}_0$.

By construction, if $k \in \mathbb{N}$ and a belief in B_i^k assigns, at information set $h \in H_i$, a non-infinitesimal probability to an opponent's strategy s_j , then s_j must be in $S_j^k(h)$, and hence there must be a belief in B_j^{k-1} for which s_j is optimal from h onwards. In other words, a belief in B_i^k believes, at every information set, that each opponent will choose rationally now and in the future. This resembles the idea of *belief in the opponents' future rationality* as formalized in Perea (2014).⁶

Note that in the definition of perfect backwards rationalizability, the beliefs for player i can always be summarized by a pair (b_i^{self}, b_i^{opp}) . Thus, we implicitly assume that player i 's belief about his own future choices is always independent from his belief about the opponents' future choices.

One important question, of course, is whether in every finite dynamic game we can find for every player at least one strategy that is perfect backwards rationalizable. The answer to this question is “yes”, as will be shown by the following theorem.

Theorem 3.1 (*Existence*). For every finite dynamic game, and every player i , there is at least one strategy for player i that is perfect backwards rationalizable.

The proof, which is constructive, can be found in Appendix B. We will finally illustrate the perfect backwards rationalizability procedure by means of an example.

Example 1 (*Illustration of perfect backwards rationalizability procedure*).

Consider the game in Fig. 2. This game starts at \emptyset , where player 1 can choose between a and b , and player 2 has a unique choice which we do not explicitly model. If player 1 chooses a the game reaches information set h_1 , where players 1 and 2 can simultaneously

⁴ A more accurate description would be “deems his own mistakes infinitely less likely than the opponents' mistakes”. However, we use “least likely” instead, as to keep the description short.

⁵ Here, and at other places in the definition, information set h need not belong to player i .

⁶ Similar ideas can be found in Penta (2015) and Baltag et al. (2009).




Fig. 2. Illustration of perfect backwards rationalizability.

choose between c and d , and between e and f , respectively. If he chooses b instead, the game reaches h_2 where players 1 and 2 can simultaneously choose between g and h , and between i and j , respectively.

In the analysis that follows we do not explicitly consider player 2’s beliefs at \emptyset as these are not relevant for our purposes. Remember that b_1^{self} denotes player 1’s cautious belief about his own strategy choice, whereas his cautious belief about player 2’s strategy choice is denoted by b_1^{opp} . Then, $b_1 = (b_1^{self}, b_1^{opp})$ constitutes player 1’s belief about his own behavior and player 2’s behavior. Let $b_1^{self}(\emptyset, a)$ be player 1’s conditional belief about his own future choice at h_1 , and $b_1^{self}(\emptyset, b)$ his conditional belief about his own future choice at h_2 . As an abbreviation, we denote by $b_1^{opp}(e) := b_1^{opp}(e, i) + b_1^{opp}(e, j)$ the probability that player 1 assigns to player 2 choosing e at h_1 , and similarly for $b_1^{opp}(f)$. Then, by definition, player 1’s expected utility of choosing a at \emptyset is

$$\begin{aligned}
 u_1(a, b_1, \emptyset) &= b_1^{self}(\emptyset, a)(a, c) \cdot b_1^{opp}(e) \cdot 1 + b_1^{self}(\emptyset, a)(a, d) \cdot b_1^{opp}(f) \cdot 2 \\
 &= 1 - b_1^{self}(\emptyset, a)(a, d) \cdot (1 - 3b_1^{opp}(f)) - b_1^{opp}(f).
 \end{aligned}
 \tag{3.1}$$

Here, we have used the fact that $b_1^{self}(\emptyset, a)(a, c) = 1 - b_1^{self}(\emptyset, a)(a, d)$, and that $b_1^{opp}(e) = 1 - b_1^{opp}(f)$. Similarly, player 1’s expected utility of choosing b at \emptyset is

$$u_1(b, b_1, \emptyset) = b_1^{self}(\emptyset, b)(b, g) = 1 - b_1^{self}(\emptyset, b)(b, h).
 \tag{3.2}$$

Initial step. Note that (b, g) is player 1’s only optimal strategy at h_2 , and hence (b, h) is a mistake. As in B_1^0 player 1 must believe in his own future rationality, we have that $v(b_1^{self}(\emptyset, b)(b, h)) > 0$ for all $b_1 \in B_1^0$.

Step 1. Clearly, $S_1^1(h_2) = \{(b, g)\}$. By construction, every $b_2 \in B_2^1$ must believe $S_1^1(h_2) = \{(b, g)\}$, and therefore $v(b_2^{opp}(h_2)(b, h)) > 0$ for all $b_2 \in B_2^1$.

Similarly, note that e is player 2’s only optimal choice at h_1 , and hence f is a mistake. We thus conclude that $S_2^1(h_1) = \{(e, i), (e, j)\}$. By definition, every $b_1 \in B_1^1$ must believe $S_2^1(h_1)$ at h_1 , and hence $v(b_1^{opp}(f)) > 0$ for all $b_1 \in B_1^1$.

Remember that $v(b_1^{self}(\emptyset, b)(b, h)) > 0$ for all $b_1 \in B_1^0$. Therefore, $st(b_1^{self}(\emptyset, b)(b, h)) = 0$. As every $b_1 \in B_1^0$ deems his own mistakes least likely, we see that

$$v(b_1^{self}(\emptyset, b)(b, h)) = v(b_1^{self}(\emptyset, b)(b, h) - st(b_1^{self}(\emptyset, b)(b, h))) > v(b_1^{opp}(f))$$

for all $b_1 \in B_1^0$, and hence in particular for all $b_1 \in B_1^1$. Together with the insight above that $v(b_1^{opp}(f)) > 0$ for all $b_1 \in B_1^1$, we conclude that

$$v(b_1^{self}(\emptyset, b)(b, h)) > v(b_1^{opp}(f)) > 0 \text{ for all } b_1 \in B_1^1.
 \tag{3.3}$$

Step 2. We have seen that $v(b_2^{opp}(h_2)(b, h)) > 0$ for all $b_2 \in B_2^1$. Hence, i is player 2’s unique optimal choice at h_2 for every belief $b_2 \in B_2^1$. As e is player 2’s unique optimal choice at h_1 for any belief, we conclude that $S_2^2(\emptyset) = \{(e, i)\}$.

We now turn to player 1’s beliefs. By combining (3.1), (3.2) and (3.3), it holds for every $b_1 \in B_1^1$ that

$$u_1(a, b_1, \emptyset) = 1 - b_1^{self}(\emptyset, a)(a, d) \cdot (1 - 3b_1^{opp}(f)) - b_1^{opp}(f) < 1 - b_1^{opp}(f) < 1 - b_1^{self}(\emptyset, b)(b, h) = u_1(b, b_1, \emptyset).$$

Hence, b is the only optimal choice for player 1 at \emptyset for every belief $b_1 \in B_1^1$. As g is the only optimal choice for player 1 after b for every $b_1 \in B_1^1$, we conclude that $S_1^2(\emptyset) = \{(b, g)\}$.

We thus see that $S_1^2(\emptyset) = \{(b, g)\}$ and $S_2^2(\emptyset) = \{(e, i)\}$. By Theorem 3.1, there is at least one strategy for player 1 and 2 that survives the procedure, and hence (b, g) and (e, i) must be the only strategies for player 1 and 2 that survive the procedure. Therefore, (b, g) and (e, i) are the unique perfect backwards rationalizable strategies for players 1 and 2 in this game.

It turns out that these are also the only quasi-perfectly rationalizable strategies in this game. However, strategy (a, c) for player 1 is perfectly rationalizable, whereas it is not perfect backwards rationalizable. The reason is that according to perfect rationalizability, player 1 is free to believe that his own mistakes are more likely than player 2’s mistakes. In particular, player 1 is free to believe that his mistake h after choosing b is much more likely than his own mistake d , and player 2’s mistake f , after choosing a . In that case, it would be optimal for player 1 to choose a at \emptyset and c at h_1 . □

4. Relation to other solution concepts

In this section we relate perfect backwards rationalizability to the concepts of perfect rationalizability, procedural quasi-perfect rationalizability and permissibility, where the first two concepts are new.

4.1. Relation to perfect rationalizability

If in the definition of perfect backwards rationalizability we drop the condition of “own mistakes being deemed least likely” in B_i^0 , then we obtain a rationalizability analogue to Selten’s (1975) perfect equilibrium which we call perfect rationalizability.

Definition 4.1 (Perfect rationalizability). (Initial step) Set $S_i^0(h) := S_i(h)$ and

$$B_i^0 := \{b_i \in B_i \mid b_i \text{ believes in his own future rationality}\}$$

for all players i and all information sets h .

(Inductive step) Let $k \in \mathbb{N}$, and suppose that $S_i^{k-1}(h)$ and B_i^{k-1} have been defined for all players i and all information sets h . Then define, for every player i and every information set h ,

$$S_i^k(h) := \{s_i \in S_i^{k-1}(h) \mid \text{there is some } b_i \in B_i^{k-1} \text{ such that } s_i(h') \text{ is locally rational for } b_i \text{ at every } h' \in H_i(s_i) \text{ weakly following } h\}$$

and

$$B_i^k := \{b_i = (b_i^{self}, b_i^{opp}) \in B_i^{k-1} \mid b_i^{opp}(h) \text{ believes } S_{-i}^k(h) \text{ for all } h \in H_i\}.$$

A strategy $s_i \in S_i$ is called perfectly rationalizable if $s_i \in S_i^k(\emptyset)$ for all $k \in \mathbb{N}_0$.

The following observation is an immediate consequence of the definitions.

Remark 4.1. Every strategy that is perfect backwards rationalizable, is also perfectly rationalizable.

This remark thus states that perfect backwards rationalizability inherits all the desirable properties from perfect equilibrium, except for the “correct beliefs” conditions that separate it from perfect rationalizability.

The other direction in the remark above is not true, as can be seen from the game Γ^2 in Fig. 1. In that game, the strategy b is perfectly rationalizable but not perfect backwards rationalizable. Indeed, if player 1 believes that his own mistake d is infinitely less likely than the opponent’s mistake f , then he must go for (a, c) .

4.2. Relation to procedural quasi-perfect rationalizability

In this subsection we propose procedural quasi-perfect rationalizability as a rationalizability analogue to van Damme’s (1984) quasi-perfect equilibrium. Like perfect backwards rationalizability, procedural quasi-perfect rationalizability is defined by iteratively eliminating strategies and beliefs from the game. The main idea that distinguishes (procedural) quasi-perfect rationalizability and quasi-perfect equilibrium from perfect backwards rationalizability is that a player believes, at each of his information sets, that his opponents will always make mistakes with some positive infinitesimal probability, but that he will not make mistakes himself. Similarly

to perfect backwards rationalizability, both (procedural) quasi-perfect rationalizability and quasi-perfect equilibrium are still based on the assumption that players deem all opponents’ strategies possible, and that a player, at each of his information sets, believes in the opponents’ future rationality.

In order to formally introduce procedural quasi-perfect rationalizability, we need the following additional notation and definitions. As before, let B_i^{opp} be the set of cautious non-standard probability distributions on the set of opponents’ strategy combinations S_{-i} . For a belief $b_i^{opp} \in B_i^{opp}$ and information set $h \in H_i$, let $b_i^{opp}(h)$ be the induced conditional probability distribution on $S_{-i}(h)$. Consider an information set $h \in H_i$ and a strategy $s_i \in S_i(h)$. By

$$u_i(s_i, b_i^{opp}(h)) := \sum_{s_{-i} \in S_{-i}(h)} b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i}))$$

we denote the *expected utility* at information set $h \in H_i$ of choosing strategy s_i under the conditional belief $b_i^{opp}(h)$. We say that the strategy $s_i \in S_i(h)$ is *globally rational* for b_i^{opp} at h if

$$u_i(s_i, b_i^{opp}(h)) \geq u_i(s'_i, b_i^{opp}(h)) \text{ for all } s'_i \in S_i(h).$$

In *procedural quasi-perfect rationalizability*, we recursively define sets of strategies $S_i^k(h)$, for all information sets h , and sets of non-standard beliefs $B_i^{opp,k}$, as follows.

Definition 4.2 (*Procedural quasi-perfect rationalizability*). (Initial step) Set $S_i^0(h) := S_i(h)$ and $B_i^{opp,0} := B_i^{opp}$ for all players i and all information sets h .

(Inductive step) Let $k \in \mathbb{N}$, and suppose that $S_i^{k-1}(h)$ and $B_i^{opp,k-1}$ have been defined for all players i and all information sets h . Then define, for every player i and every information set h ,

$$S_i^k(h) := \{s_i \in S_i^{k-1}(h) \mid \text{there is some } b_i^{opp} \in B_i^{opp,k-1} \text{ such that } s_i \text{ is globally rational for } b_i^{opp} \text{ at every } h' \in H_i(s_i) \text{ weakly following } h\}$$

and

$$B_i^{opp,k} := \{b_i \in B_i^{opp,k-1} \mid b_i^{opp}(h) \text{ believes } S_{-i}^k(h) \text{ for all } h \in H_i\}.$$

A strategy $s_i \in S_i$ is procedurally quasi-perfectly rationalizable if $s_i \in S_i^k(\emptyset)$ for all $k \in \mathbb{N}_0$.

Similar to perfect backwards rationalizability, also procedural quasi-perfect rationalizability embodies the idea of *belief in the opponents’ future rationality* as proposed by Perea (2014). Indeed, if a belief in $B_i^{opp,k}$ assigns, at information set $h \in H_i$, a non-infinitesimal probability to an opponent’s strategy s_j , then s_j must be in $S_j^k(h)$, and hence there must be some belief in $B_j^{opp,k-1}$ for which s_j is optimal from h onwards. The crucial difference between procedural quasi-perfect rationalizability and perfect backwards rationalizability is that in the latter concept, a player also deems possible future mistakes by himself, whereas in the first concept a player only takes into account mistakes by his opponents. However – and that is crucial – under the latter concept the player deems his own mistakes infinitely less likely than the opponents’ mistakes. Due to this last property, it can be shown that every perfect backwards rationalizable strategy is also procedurally quasi-perfectly rationalizable.

Theorem 4.1 (*Relation with procedural quasi-perfect rationalizability*). *Every perfect backwards rationalizable strategy is procedurally quasi-perfectly rationalizable.*

The other direction is not true. Indeed, in game Γ^1 from Fig. 1 the strategy (a, c) is procedurally quasi-perfectly rationalizable but not perfect backwards rationalizable.

The proof of Theorem 4.1 can be found in Appendix C. What makes the proof challenging is that perfect backwards rationalizability and procedural quasi-perfect rationalizability are defined in fundamentally different ways: In perfect backwards rationalizability a player holds beliefs about his own future choices, and optimality of a strategy is defined *locally*, on a choice-by-choice basis. That is, optimality requires that at every information set the prescribed choice is locally optimal, given the player’s belief about his own future choices and given his belief about the opponents’ strategies. In contrast, procedural quasi-perfect rationalizability does not involve beliefs about the player’s own future choices, and optimality is defined *globally*. That is, optimality requires that at every information set the player’s strategy is optimal, given his belief about the opponents’ strategies. A key step in the proof is to show that, given a fixed belief, a sequence of locally optimal *choices* in the perfect backwards rationalizability concept always yields a globally optimal *strategy* in the procedural quasi-perfect rationalizability concept. See Corollary C.1 in Appendix C.

It is easily seen that procedural quasi-perfect rationalizability is a refinement of the backwards rationalizability procedure in Perea (2014) and Penta (2015). Indeed, if in the definition of procedural quasi-perfect rationalizability we replace B_i^{opp} by the set of standard conditional belief vectors (with zero non-standard part) satisfying Bayesian updating, then we obtain exactly the definition of backwards rationalizability. Note that every cautious non-standard belief $b_i^{opp} \in B_i^{opp}$ naturally induces a standard conditional belief vector satisfying Bayesian updating, by taking at every information set $h \in H_i$ the standard part of $b_i^{opp}(h)$. Moreover, if a

strategy s_i is globally rational at $h \in H_i(s_i)$ for b_i^{opp} , then in particular it is optimal at h for the standard part of $b_i^{opp}(h)$. That is, global rationality at h for b_i^{opp} implies global rationality for the induced standard conditional belief vector. These two insights together imply the following result.

Remark 4.2. Every strategy that survives the procedural quasi-perfect rationalizability procedure is backwards rationalizable.

The key difference between the two procedures is thus that the backwards rationalizability procedure does not impose cautious reasoning, as a player, at each of his information sets, is free to assign probability 0 to certain opponents' strategies. If we combine Remark 4.2 with Theorem 4.1, we arrive at the following result.

Remark 4.3. Every perfect backwards rationalizable strategy is backwards rationalizable.

Together with Remark 4.1 this justifies the name *perfect backwards rationalizability*, as it may be viewed as a perfection-based refinement of backwards rationalizability.

Perea (2014) has shown that in every game with perfect information without relevant ties, the only strategies that survive the backwards rationalizability procedure are the backward induction strategies. In the light of Remark 4.3 it thus follows that in every such game, the only perfect backwards rationalizable strategies are the backward induction strategies.

Since we have seen in the previous section that perfect backwards rationalizability is a refinement of perfect rationalizability, it follows from Theorem 4.1 that perfect backwards rationalizability refines both perfect rationalizability and procedural quasi-perfect rationalizability. Hence, it inherits all the desirable properties that perfect rationalizability and procedural quasi-perfect rationalizability display. Yet, it adds the requirement that a player deems his own future mistakes infinitely less likely than his opponents' mistakes, without completely discarding his own future mistakes as procedural quasi-perfect rationalizability does.

Corollary 4.1 (*Relation with perfect and procedural quasi-perfect rationalizability*). *Every perfect backwards rationalizable strategy is both perfectly rationalizable and procedurally quasi-perfectly rationalizable.*

In particular, perfect backwards rationalizability rules out both strategies containing weakly dominated actions in the agent normal form, as well as weakly dominated strategies in the normal form. This follows from the fact that perfect rationalizability avoids strategies containing weakly dominated actions in the agent normal form, and procedural quasi-perfect rationalizability avoids weakly dominated strategies in the normal form.

In each of the examples we have seen so far, the other direction of Corollary 4.1 was also true. Indeed, in each of those examples every strategy that was both perfectly rationalizable and procedurally quasi-perfectly rationalizable was also perfect backwards rationalizable. It is easily seen that this direction is always true in every game with perfect information and without relevant ties, as in such games the three concepts above all uniquely select the backward induction strategies for the players. Whether this direction is still true for games with perfect information and relevant ties is still an open question to us.

However, as the following example will show, the opposite direction of the corollary is not generally true for games with imperfect information. There are games where a strategy is both perfectly rationalizable and procedurally quasi-perfectly rationalizable, but not perfect backwards rationalizable.

Example 2 (*Combining perfect and procedural quasi-perfect rationalizability does not lead to perfect backwards rationalizability*).

Consider the game in Fig. 3. Then, strategy g for player 2 is both perfectly rationalizable and procedurally quasi-perfectly rationalizable, but not perfect backwards rationalizable. To see that g is perfectly rationalizable, note first that strategy b is perfectly rationalizable for player 1 if he believes that his own mistake (c, e) is much more likely than player 2's mistake i . Hence, under perfect rationalizability, player 2 may assign at his information set a high probability to player 1 choosing b , which makes g optimal for player 2.

To see that g is procedurally quasi-perfectly rationalizable, note that under procedural quasi-perfect rationalizability player 1 believes that he will not make mistakes himself, and hence strategy (a, c, f) will be among his optimal strategies. Therefore, player 2 may assign at his information set a high probability to player 1 choosing (a, c, f) , which makes g optimal for player 2.

Under perfect backwards rationalizability, however, player 1 believes that he will make the mistake (c, e) with positive probability, but he deems the probability of player 2 making the mistake i much higher. As such, the only perfect backwards rationalizable strategy for player 1 is (a, d) . Hence, player 2 must at his information set assign a high probability to player 1 choosing (a, d) , which implies that player 2 must choose h . That is, h is the only perfect backwards rationalizable strategy for player 2. In particular, g is not perfect backwards rationalizable.

4.3. Relation to permissibility

Suppose next that we apply the definition of perfect backwards rationalizability to a *one-shot game*, in which all players simultaneously make one choice at the initial history \emptyset , after which the game ends. In that case, we obtain the concept of *permissibility* (Brandenburger (1992), Börgers (1994)). To see this, note that in a one-shot game there are no future choices, and hence the set B_i would only contain beliefs about opponents' strategies. That is, $B_i = B_i^{opp}$ contains all cautious beliefs about the opponents' strategy




Fig. 3. Combining perfect and procedural quasi-perfect rationalizability does not lead to perfect backwards rationalizability.

combinations. Moreover, the conditions of *belief in own future rationality* and *deeming own mistakes least likely* are vacuous since there are no future choices. Finally, S_i would simply be the set of choices for player i at \emptyset , and we can write S_i^k instead of $S_i^k(h)$ as there is only one information set in the game. The procedure obtained would be as follows.

Definition 4.3 (One-shot perfect backwards rationalizability). (Initial step) Set $S_i^0 := S_i$ and $B_i^0 := B_i^{opp}$ for all players i . (Inductive step) Let $k \in \mathbb{N}$, and suppose that S_i^{k-1} and B_i^{k-1} have been defined for all players i . Then define, for every player i ,

$$S_i^k := \{s_i \in S_i^{k-1} \mid \text{there is some } b_i \in B_i^{k-1} \text{ such that } s_i \text{ is rational for } b_i\}$$

and

$$B_i^k := \{b_i \in B_i^{k-1} \mid b_i \text{ believes } S_{-i}^k\}.$$

A strategy $s_i \in S_i$ is called one-shot perfect backwards rationalizable if $s_i \in S_i^k$ for all $k \in \mathbb{N}_0$.

Here, we say that s_i is rational for b_i if $u_i(s_i, b_i, \emptyset) \geq u_i(s'_i, b_i, \emptyset)$ for all $s'_i \in S_i$. If in the procedure above we replace cautious non-standard beliefs by cautious lexicographic beliefs,⁷ then we obtain precisely the *permissibility* concept defined in Brandenburger (1992). Indeed, the key condition above that b_i believes S_{-i}^k , when translated into lexicographic beliefs, would correspond to requiring that the lexicographic belief b_i assigns first-order probability 1 to S_{-i}^k , which is precisely the driving condition in Brandenburger’s definition. We thus arrive at the following conclusion.

Remark 4.4. In a one-shot game, perfect backwards rationalizability is equivalent to permissibility.

Brandenburger (1992) has shown that permissibility is equivalent to the *Dekel-Fudenberg procedure* (Dekel and Fudenberg (1990)), in which we start by one round of elimination of weakly dominated strategies, followed by the iterated elimination of strictly dominated strategies. In view of Remark 4.4, it follows that perfect backwards rationalizability is equivalent to the Dekel-Fudenberg procedure when applied to a one-shot game. The same is true for perfect rationalizability and (procedural) quasi-perfect rationalizability.

5. Discussion

In this section we discuss why we use non-standard probabilities, instead of lexicographic probability systems, to define the concept of perfect backwards rationalizability, and how the concept relates to *quasi-perfect rationalizability* and *perfect quasi-perfect equilibrium*.

⁷ A *lexicographic belief* (or *lexicographic probability system*) for player i is a finite sequence of beliefs $b_i = (b_i^1; b_i^2; \dots; b_i^K)$, where $b_i^k \in \Delta(S_{-i})$ for all $k \in \{1, \dots, K\}$. The lexicographic belief b_i is *cautious* if for every $s_{-i} \in S_{-i}$ there is some $k \in \{1, \dots, K\}$ with $b_i^k(s_{-i}) > 0$.

5.1. Lexicographic probability systems

Recall that we have been using *non-standard probability distributions* (Robinson (1973), Hammond (1994) and Halpern (2010)) with infinitesimals to model beliefs about mistakes. This leaves the question why we did not opt for *lexicographic probability systems* (Blume et al. (1991)) to model such cautious beliefs, as is common nowadays in epistemic game theory. The reason is that non-standard probabilities allow us to take the product of two beliefs in a very easy way, whereas this is considerably more difficult (although possible) with lexicographic probability systems. This advantage is important, since in the concept of perfect backwards rationalizability, a player holds, at each of his information sets, both (a) a cautious belief about his *own* future choices, and (b) a cautious belief about the *opponents'* strategies. To determine which choice is optimal for the player at that information set, we must take the “product” of the belief about his own choices and the belief about the opponents' choices. Taking such products is rather complicated for lexicographic probability distributions, whereas it comes for free when using non-standard probability distributions.⁸

5.2. Relation to quasi-perfect rationalizability

Asheim and Perea (2005)'s definition of quasi-perfect rationalizability differs both methodologically and behaviorally from procedural quasi-perfect rationalizability. Instead of using a procedure that recursively eliminates strategies and beliefs from the game, Asheim and Perea use *belief hierarchies* as a primitive notion to define quasi-perfect rationalizability. That is, Asheim and Perea do not only consider first-order beliefs about the opponents' strategies, as we do in procedural quasi-perfect rationalizability, but also explore the players' *second-order* beliefs about the opponents' beliefs about the strategies of others, and *higher-order beliefs* as well. Asheim and Perea encode such belief hierarchies by means of epistemic models with types and lexicographic beliefs, and impose epistemic conditions on such belief hierarchies which give rise to their definition of quasi-perfect rationalizability.

In Appendix D we formally define quasi-perfect rationalizability and explore, in detail, the relation between procedural quasi-perfect rationalizability and quasi-perfect rationalizability. We show that every strategy that is quasi-perfectly rationalizable is also procedurally quasi-perfectly rationalizable, but not *vice versa*. Hence, quasi-perfect rationalizability is stronger than procedural quasi-perfect rationalizability. Intuitively, the key difference is the following: According to procedural quasi-perfect rationalizability, if a player i at information set h deems an opponent's strategy s_j most plausible, then there is a belief b_j that survives all rounds and for which the strategy s_j is optimal from h onwards. Quasi-perfect rationalizability requires more: If player i , at information set h , deems an opponent's belief b_j possible, and at (a possibly different) information set h' deems the opponent's strategy s_j most plausible given the opponent's belief b_j , then the strategy s_j must be optimal for b_j from h' onwards. Here, when we say that player i deems strategy s_j most plausible at h , we mean that there is no other strategy $s'_j \in S_j(h)$ that b_i deems infinitely more likely than s_j . Hence, according to quasi-perfect rationalizability, the opponent's belief b_j that player i assigns to his opponent at information set h is not only used to justify his behavior from h onwards, but also to form his belief about j 's behavior at information sets that do not follow h . In that sense, quasi-perfect rationalizability imposes restrictions that go beyond belief in the opponents' future rationality. In contrast, procedural quasi-perfect rationalizability only imposes rationality restrictions that are in line with belief in the opponents' future rationality.

Recall from Theorem 4.1 that every perfect backwards rationalizable strategy is also procedurally quasi-perfect rationalizable. In Appendix D we show that this is no longer true if we replace procedural quasi-perfect rationalizability by quasi-perfect rationalizability. Indeed, we provide a counterexample where some strategy is perfect backwards rationalizable, but not quasi-perfectly rationalizable. Apparently, the extra conditions that quasi-perfect rationalizability imposes relative to procedural quasi-perfect rationalizability, as discussed at the beginning of this section, are not shared by perfect backwards rationalizability. At the same time, quasi-perfect rationalizability is not a refinement of perfect backwards rationalizability, as can be seen from the game Γ^1 in Fig. 1. In that game, strategy (a, c) is quasi-perfectly rationalizable, but not perfect backwards rationalizable.

5.3. Relation to perfect quasi-perfect equilibrium

The concept of *perfect quasi-perfect equilibrium* developed in Blume and Meier (2019) may be viewed as an equilibrium counterpart to perfect backwards rationalizability. Unlike our concept, it (implicitly) imposes a correct beliefs assumption, requiring that every player believes that his opponents are correct about the beliefs he holds.

The main other difference lies in the way beliefs about mistakes are modeled. Instead of using non-standard probability distributions, Blume and Meier (2019) simulate (beliefs about) mistakes by trembles – that is, sequences of full-support probability distributions that converge to the strategy profile under consideration. When evaluating the strategy of player i , Blume and Meier (2019) also consider trembles in player i 's own strategy, comparable to how perfect equilibrium (Selten (1975)) proceeds. Similarly to our approach, the key condition in perfect quasi-perfect equilibrium is that player i deems the trembles in his own strategy much smaller, in fact *infinitely smaller*, than the trembles in the opponents' strategies.

Recall from Corollary 4.1 that perfect backwards rationalizability constitutes a refinement of both perfect rationalizability and procedural quasi-perfect rationalizability. Interestingly, this result does not have an equilibrium counterpart. In Blume and Meier

⁸ An alternative approach would be to consider lexicographic beliefs about the combinations of own continuation strategies and opponents' strategies, and requiring this belief to be independent across these two components. Deeming your own mistakes least likely would then correspond to additional restrictions on such lexicographic beliefs.

(2019) it is shown that perfect quasi-perfect equilibrium is a refinement of quasi-perfect equilibrium but not of perfect equilibrium. Even more, they show that there are dynamic games where the sets of quasi-perfect equilibria and perfect equilibria are disjoint, rendering Corollary 4.1 impossible in an equilibrium framework. Apparently, the correct beliefs assumption which is implicit in equilibrium concepts makes the logic behind perfect equilibrium and quasi-perfect equilibrium incompatible. However, in view of Corollary 4.1, both logics become compatible once we drop the correct beliefs assumption.

6. Concluding remarks

This paper presents a new rationalizability concept for dynamic games, based on three principles: (a) a player always believes in his own future rationality and in the opponents’ future rationality, (b) a player always believes that he himself, and his opponents, will make mistakes with positive infinitesimal probability, and (c) a player deems his own mistakes infinitely less likely than those of his opponents. The concept may open the door to variations, or extensions, across different dimensions.

First, it could be extended – probably without major difficulties – to dynamic games with incomplete information. Second, condition (a), which guarantees that the concept has a backward induction flavor, could be replaced, for instance, by *strong belief in the opponents’ rationality* (Battigalli and Siniscalchi (2002)), turning it into a forward induction concept. Third, as discussed in the introduction, the concept could be weakened if condition (c) is replaced by the weaker requirement that a player deems his own mistakes less likely, but not necessarily infinitely less likely, than those of his opponents.

Declaration of competing interest

None.

Appendix A. Non-standard probabilities

A.1. Non-standard numbers

Recall that the field of non-standard numbers $\mathbb{R}(\epsilon)$ contains all numbers a that can be written as

$$a = \frac{a_0 + a_1\epsilon + a_2\epsilon^2 + \dots + a_K\epsilon^K}{b_0 + b_1\epsilon + b_2\epsilon^2 + \dots + b_K\epsilon^K}, \tag{A.1}$$

where $a_k, b_k \in \mathbb{R}$ for all $k \in \{0, \dots, K\}$, $b_k \neq 0$ for some $k \in \{0, \dots, K\}$, and where either $a_0 \neq 0$ or $b_0 \neq 0$. We call the non-standard number $a \in \mathbb{R}(\epsilon)$ *finite* if $b_0 \neq 0$. We now show that every finite non-standard number can be written as a (possibly infinite) polynomial in ϵ . Since $b_0 \neq 0$, we can write the denominator in (A.1) as

$$b_0 \left(1 + \frac{b_1}{b_0}\epsilon + \frac{b_2}{b_0}\epsilon^2 + \dots + \frac{b_K}{b_0}\epsilon^K \right).$$

Moreover, by the property of ϵ we know that $\left| \frac{b_k}{b_0}\epsilon^k \right| < \left(\frac{1}{2}\right)^k$ for every $k \in \{1, \dots, K\}$, and hence

$$\left| \frac{b_1}{b_0}\epsilon + \frac{b_2}{b_0}\epsilon^2 + \dots + \frac{b_K}{b_0}\epsilon^K \right| \leq \sum_{k=1}^K \left| \frac{b_k}{b_0}\epsilon^k \right| < \sum_{k=1}^K \left(\frac{1}{2}\right)^k < 1.$$

But then, by the formula for geometric series it immediately follows that

$$\left(1 + \frac{b_1}{b_0}\epsilon + \frac{b_2}{b_0}\epsilon^2 + \dots + \frac{b_K}{b_0}\epsilon^K \right)^{-1} = 1 + \sum_{m=1}^{\infty} (-1)^m \left(\frac{b_1}{b_0}\epsilon + \frac{b_2}{b_0}\epsilon^2 + \dots + \frac{b_K}{b_0}\epsilon^K \right)^m.$$

Combining this with (A.1) then yields

$$a = \frac{1}{b_0} (a_0 + a_1\epsilon + a_2\epsilon^2 + \dots + a_K\epsilon^K) \left(1 + \sum_{m=1}^{\infty} (-1)^m \left(\frac{b_1}{b_0}\epsilon + \frac{b_2}{b_0}\epsilon^2 + \dots + \frac{b_K}{b_0}\epsilon^K \right)^m \right)$$

which is a power series in ϵ . We thus conclude that every *finite* number $a \in \mathbb{R}(\epsilon)$ can be written as

$$a = a_0 + a_1\epsilon + a_2\epsilon^2 + \dots, \tag{A.2}$$

where $a_k \in \mathbb{R}$ for every $k \in \mathbb{N}_0$. We call this a *power series representation* of the number a .⁹ Below, we will show that this power series representation is unique.

⁹ If the number a is not finite, then it will have a power series representation of the form $a_0\epsilon^{-k} + a_1\epsilon^{-k+1} + \dots$ starting with a term having a negative exponent. This would correspond to a *Laurent series*. For finite numbers, the power series representation is a *Taylor series*. Since we are only interested in finite numbers, we do not need Laurent series in this paper.

A.2. Properties of non-standard numbers

Consider a finite non-standard number $a = a_0 + a_1\varepsilon + a_2\varepsilon^2 + \dots$. Then, we say that $a = 0$ if $a_k = 0$ for all $k \in \mathbb{N}_0$. This immediately implies that the power series representation of any finite number a is unique. Indeed, suppose that

$$a = a_0 + a_1\varepsilon + a_2\varepsilon^2 + \dots = b_0 + b_1\varepsilon + b_2\varepsilon^2 + \dots$$

Then,

$$(a_0 - b_0) + (a_1 - b_1)\varepsilon + (a_2 - b_2)\varepsilon^2 + \dots = 0,$$

which means that $a_k = b_k$ for all $k \in \mathbb{N}_0$. We can therefore refer to (A.2) as *the* power series representation of the non-standard number a .

If $a \neq 0$, we denote by $v(a)$ the smallest index k for which $a_k \neq 0$, and call it the *valuation* of the number a . We set $v(0) := \infty$. For a finite number $a \neq 0$, we write $a > 0$ if $a_{v(a)} > 0$, and $a < 0$ if $a_{v(a)} < 0$. For two finite numbers a, b , we say that $a > b$ if $a - b > 0$. Moreover, we say that a is of *infinitely smaller size* than b if $v(a) > v(b)$. We use the term “infinitely smaller size” rather than the more familiar “infinitely smaller” because we also apply it to negative numbers. For instance, ε is of infinitely smaller size than -1 , although ε is not smaller than -1 . Note that, by definition, 0 is of infinitely smaller size than any finite non-zero non-standard number.

For a finite non-standard number a with power series representation (A.2), and a given $k \in \mathbb{N}_0$, we call

$$\text{trunc}^k(a) = a_0 + a_1\varepsilon + \dots + a_k\varepsilon^k$$

the k -th order truncation of a . The 0 -th order truncation $a_0 \in \mathbb{R}$ is also called the *standard part* of a , and is denoted by $st(a)$. Hence, $st(a)$ is the unique real number that is closest to a .

It turns out that the k -th order truncation of a will either be zero, or have the same sign as a . This property will be important for our proofs.

Lemma A.1 (*Truncation has the same sign*). Consider a finite number $a \in \mathbb{R}(\varepsilon)$ with $a \geq 0$. Then, $\text{trunc}^k(a) \geq 0$ for every $k \in \mathbb{N}_0$.

Proof. Suppose first that $a = 0$. Then, $a_k = 0$ for all $k \in \mathbb{N}_0$, and hence $\text{trunc}^k(a) = 0$ for all $k \in \mathbb{N}_0$. Assume next that $a > 0$. Then, $a_{v(a)} > 0$. If $k < v(a)$, then $\text{trunc}^k(a) = 0$. If $k \geq v(a)$, then $v(\text{trunc}^k(a)) = v(a)$, and hence $\text{trunc}^k(a) > 0$. \square

In the following subsection we will use the property above to investigate non-standard probability distributions.

A.3. Non-standard probability distributions

Consider a finite set X . A *non-standard probability distribution* on X is a function $p : X \rightarrow \mathbb{R}(\varepsilon)$ such that $p(x) \geq 0$ for all $x \in X$ and $\sum_{x \in X} p(x) = 1$. By $\Delta^{ns}(X)$ we denote the set of non-standard probability distributions on X . Such non-standard probability distributions will often be interpreted as *beliefs*. We therefore use the terms “non-standard probability distribution” and “belief” interchangeably in this paper. For two elements x and y in X , we say that p deems x *infinitely more likely than* y if $p(y)$ is of infinitely smaller size than $p(x)$.

Consider a non-standard probability distribution p on X . For a subset $Y \subseteq X$ with $\sum_{x \in Y} p(x) > 0$, the *conditional probability distribution* on Y induced by p is the non-standard probability distribution p_Y on Y given by

$$p_Y(x) := \frac{p(x)}{\sum_{y \in Y} p(y)}$$

for every $x \in Y$.

We say that p is *cautious* on X if $p(x) > 0$ for all $x \in X$, such that conditional probability distributions can be formed for every subset $Y \subseteq X$. We call p a *standard probability distribution* on X if $p(x) \in \mathbb{R}$ for all $x \in X$, and the set of standard probability distributions on X is denoted by $\Delta(X)$. A *standard zero-sum distribution* on X is a function $f : X \rightarrow \mathbb{R}$ with $\sum_{x \in X} f(x) = 0$.

Consider a non-standard probability distribution p on X . From above we know that every probability $p(x)$ has a unique power series representation

$$p(x) = p_0(x) + p_1(x)\varepsilon + p_2(x)\varepsilon^2 + \dots,$$

where $p_k(x) \in \mathbb{R}$ for every $k \in \mathbb{N}_0$. As $\sum_{x \in X} p(x) = 1$, it follows that

$$\left(\sum_{x \in X} p_0(x) - 1 \right) + \varepsilon \left(\sum_{x \in X} p_1(x) \right) + \varepsilon^2 \left(\sum_{x \in X} p_2(x) \right) + \dots = 0.$$

Hence,

$$\sum_{x \in X} p_0(x) = 1 \text{ and } \sum_{x \in X} p_k(x) = 0 \text{ for all } k \in \mathbb{N}.$$

Moreover, since $p(x) \geq 0$ for every $x \in X$, it follows by Lemma A.1 that $p_0(x) = \text{trunc}^0(p(x)) \geq 0$ for every $x \in X$. Thus, we conclude that $p_0 := (p_0(x))_{x \in X}$ is a standard probability distribution in $\Delta(X)$, and that $p_k := (p_k(x))_{x \in X}$ is a standard zero-sum distribution on X .

As such, every non-standard probability distribution p on X can uniquely be written as

$$p = p_0 + \varepsilon p_1 + \varepsilon^2 p_2 + \dots \tag{A.3}$$

where $p_0 \in \Delta(X)$ is a standard probability distribution on X , and p_k is a standard zero-sum distribution on X for every $k \in \mathbb{N}$. We call this the (unique) *power series representation* of the non-standard probability distribution p . This representation will be important for our game-theoretic analysis later on.

By $st(p) := p_0$ we denote the standard part of the non-standard probability distribution p . For a subset $Y \subseteq X$ we say that p *believes* Y if $st(\sum_{x \in Y} p(x)) = 1$. By the representation (A.3), this thus means that $\sum_{x \in Y} p_0(x) = 1$.

Consider an element $x \in X$. As $p(x) \geq 0$, it follows from (A.3) and Lemma A.1 that $\text{trunc}^k(p(x)) = p_0(x) + \varepsilon p_1(x) + \dots + \varepsilon^k p_k(x) \geq 0$ for every $k \in \mathbb{N}_0$. Hence, $p_0 + \varepsilon p_1 + \dots + \varepsilon^k p_k$ is a non-standard probability distribution on X as well.

Suppose that p is cautious on X , and let $k \in \mathbb{N}_0$ be the minimal index such that the truncated non-standard probability distribution $p_0 + \varepsilon p_1 + \dots + \varepsilon^k p_k$ is cautious on X . Then, $p_0 + \varepsilon p_1 + \dots + \varepsilon^k p_k$ is called the *minimal cautious truncation* of p on X .

Appendix B. Proof of Theorem 3.1

Remember that B_i^k denotes the set of belief pairs $(b_i^{\text{self}}, b_i^{\text{opp}})$ for player i that survives round k of the procedure. Similarly, $S_i^k(h)$ is the set of strategies for player i that survive round k of the procedure at information set h . By construction, $B_i^k \subseteq B_i^{k-1}$ and $S_i^k(h) \subseteq S_i^{k-1}(h)$ for all $k \geq 1$. Since the collection of information sets is finite, and the set of strategies $S_i(h)$ is finite for every player i and every information set h , the procedure must terminate within finitely many steps. To prove the existence of perfect backwards rationalizable strategies, it is therefore sufficient to show that B_i^k and $S_i^k(h)$ are always non-empty for every player i , every information set h and every $k \geq 0$. We prove so by induction on k .

For $k = 0$ we have that $S_i^0(h) = S_i(h)$, and hence $S_i^0(h)$ is non-empty. To prove that B_i^0 is non-empty, we must show that there is a belief pair $(b_i^{\text{self}}, b_i^{\text{opp}})$ for player i that believes in his own future rationality and deems his own mistakes least likely.

Take an arbitrary cautious non-standard probability distribution b_i^{opp} on the set S_{-i} of opponents' strategy combinations. For every opponents' strategy combination, let $v(b_i^{\text{opp}}(s_{-i}))$ be the valuation of the probability $b_i^{\text{opp}}(s_{-i})$, as defined in Section 5.2. That is, if $b_i^{\text{opp}}(s_{-i}) = a_0 + a_1 \varepsilon + a_2 \varepsilon^2 + \dots$, then $v(b_i^{\text{opp}}(s_{-i}))$ is the smallest number m such that $a_m \neq 0$. Let $v := \max\{v(b_i^{\text{opp}}(s_{-i})) \mid s_{-i} \in S_{-i}\}$ be the maximal valuation of $b_i^{\text{opp}}(s_{-i})$ across all opponents' strategy combinations $s_{-i} \in S_{-i}$.

We now define the cautious non-standard probability distribution b_i^{self} on the set S_i of i 's own strategies by a backward induction construction, as follows. For every $m \geq 0$, let H_i^m be the collection of information sets in H_i that are followed by at most m consecutive information sets in H_i .

We start by considering all information sets in H_i^0 , that is, player i information sets that are not followed by any other player i information set. Consider an information set $h \in H_i^0$, and let $c_i^*(h)$ be an optimal choice for player i at h given the conditional belief $b_i^{\text{opp}}(h)$. Let σ_{ih} be the cautious non-standard probability distribution on the set of available choices $C_i(h)$ given by

$$\sigma_{ih}(c_i) := \begin{cases} 1 - (|C_i(h)| - 1) \cdot \varepsilon^{v+1}, & \text{if } c_i = c_i^*(h) \\ \varepsilon^{v+1}, & \text{if } c_i \neq c_i^*(h) \end{cases} \tag{B.1}$$

Now let $m \in \mathbb{N}$ and consider some $h \in H_i^m$. Suppose that the choice $c_i^*(h')$ and the cautious non-standard probability distribution $\sigma_{ih'}$ have been defined at all $h' \in H_i^l$ where $l \leq m - 1$. In particular, $c_i^*(h')$ and $\sigma_{ih'}$ have been defined for all information sets $h' \in H_i$ following h . For every choice $c_i \in C_i(h)$, let $u_i(c_i, ((\sigma_{ih'})_{h' \in H_i: h' > h}, b_i^{\text{opp}}), h)$ be the expected utility of making choice c_i at h , given the conditional belief $b_i^{\text{opp}}(h)$ about the opponents' strategy combinations, and given the non-standard probability distributions $\sigma_{ih'}$ on i 's own choices at h' for every $h' \in H_i$ that follows h . Above, we have used the expression " $h' > h$ " as a shortcut for " h' follows h ". Let $c_i^*(h)$ be an optimal choice for player i at h , that is,

$$u_i(c_i^*(h), ((\sigma_{ih'})_{h' \in H_i: h' > h}, b_i^{\text{opp}}), h) \geq u_i(c_i, ((\sigma_{ih'})_{h' \in H_i: h' > h}, b_i^{\text{opp}}), h) \text{ for all } c_i \in C_i(h). \tag{B.2}$$

Moreover, let σ_{ih} be the cautious non-standard probability distribution on the set of available choices $C_i(h)$ given by

$$\sigma_{ih}(c_i) := \begin{cases} 1 - (|C_i(h)| - 1) \cdot \varepsilon^{v+1}, & \text{if } c_i = c_i^*(h) \\ \varepsilon^{v+1}, & \text{if } c_i \neq c_i^*(h) \end{cases} \tag{B.3}$$

By induction on m we have thus defined, for every information set $h \in H_i$, the cautious non-standard probability distribution σ_{ih} on $C_i(h)$.

Let b_i^{self} be the cautious non-standard probability distribution on i 's own strategies given by

$$b_i^{\text{self}}(s_i) := \prod_{h \in H_i(s_i)} \sigma_{ih}(s_i(h)) \text{ for every } s_i \in S_i. \tag{B.4}$$

We will now show that $b_i = (b_i^{self}, b_i^{opp})$ believes in his own future rationality and deems his own mistakes least likely.

To prove that b_i believes in his own future rationality, we must show that $b_i^{self}(h, c_i)$ believes $S_i^{rat}(b_i, h)$ for every $h \in H_i$ and $c_i \in C_i(h)$. Take some $h \in H_i$ and $c_i \in C_i(h)$. By (B.4) we conclude that

$$u_i(c_i, b_i, h) = u_i(c_i, ((\sigma_{ih'})_{h' \in H_i: h' > h}, b_i^{opp}), h). \tag{B.5}$$

By (B.5) and (B.2) it then follows that

$$u_i(c_i^*(h), b_i, h) \geq u_i(c_i, b_i, h) \text{ for all } c_i \in C_i(h). \tag{B.6}$$

Let $s_i^*(h, c_i)$ be the unique strategy in $S_i(h, c_i)$ such that $s_i^*(h, c_i)$ prescribes the optimal choice $c_i^*(h')$ at every $h' \in H_i(s_i^*(h, c_i))$ not weakly preceding h . Hence, in particular, $s_i^*(h, c_i)$ prescribes the optimal choice $c_i^*(h')$ at every $h' \in H_i(s_i^*(h, c_i))$ following h . Then, by (B.6) applied to every $h' \in H_i(s_i^*(h, c_i))$ following h , we conclude that $s_i^*(h, c_i) \in S_i^{rat}(b_i, h)$.

Moreover, by (B.4) and (B.3) we conclude that the standard part of the conditional non-standard probability distribution $b_i^{self}(h, c_i)$ assigns probability 1 to $s_i^*(h, c_i) \in S_i^{rat}(b_i, h)$. This implies that $b_i^{self}(h, c_i)$ believes $S_i^{rat}(b_i, h)$. As this is true for every $h \in H_i$ and $c_i \in C_i(h)$, we conclude that b_i believes in his own future rationality.

We next prove that $b_i = (b_i^{self}, b_i^{opp})$ deems his own mistakes least likely. Consider an information set $h \in H_i$ and a choice $c_i \in C_i(h)$. From (B.4) and (B.3) we see that for every strategy $s_i \in S_i(h, c_i)$, the infinitesimal mistake part $b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i)(s_i))$ has a valuation which is at least $v + 1$. On the other hand, the non-standard probability $b_i^{opp}(s_{-i})$ has a valuation of at most v for every opponents' strategy combination s_{-i} , by definition of v . We thus conclude that $b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i)(s_i))$ is of infinitely smaller size than $b_i^{opp}(s_{-i})$. As this holds for every information set $h \in H_i$, choice $c_i \in C_i(h)$, strategy $s_i \in S_i(h, c_i)$ and opponents' strategy combination $s_{-i} \in S_{-i}$, we know that $b_i = (b_i^{self}, b_i^{opp})$ deems his own mistakes least likely.

Overall, we have thus constructed a belief $b_i = (b_i^{self}, b_i^{opp})$ that believes in his own future rationality and deems his own mistakes least likely. Hence, by definition, $b_i \in B_i^0$, which implies that B_i^0 is non-empty.

Now, let $k \in \mathbb{N}$ and assume that $S_i^{k-1}(h)$ and B_i^{k-1} are non-empty for every player i and every information set h . Consider some player i and information set h . We will show that $S_i^k(h)$ and B_i^k are non-empty.

To show that $S_i^k(h)$ is non-empty, take some $b_i = (b_i^{self}, b_i^{opp})$ in B_i^{k-1} . This is possible since we assume that B_i^{k-1} is non-empty. At every information set $h' \in H_i$, let $c_i[h']$ be a locally rationally choice for $b_i = (b_i^{self}, b_i^{opp})$ at h' . Let s_i be a strategy in $S_i(h)$ such that $s_i(h') = c_i[h']$ for every $h' \in H_i(s_i)$ weakly following h . Then, by construction, $s_i \in S_i^k(h)$, and hence $S_i^k(h)$ is non-empty.

We will next construct a belief $b_i = (b_i^{self}, b_i^{opp})$ in B_i^k . We start by defining b_i^{opp} . That is, we must find a belief b_i^{opp} such that $b_i^{opp}(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$. For every opponent $j \neq i$, let b_j be an arbitrary belief pair in B_j^{k-1} . This is possible since we assume that B_j^{k-1} is non-empty. For every information set $h \in H_i$ let $s_{-i}[h] = (s_j[h])_{j \neq i}$ be an opponents' strategy combination in $S_{-i}(h)$ with the following property: For every opponent $j \neq i$ and every information set $h' \in H_j(s_j)$ that does not precede h , the choice $(s_j[h])(h')$ is locally rational for b_j at h' . Clearly, such a strategy $s_j[h]$ can always be found. Then, by construction, $s_j[h] \in S_j^k(h')$ for every h' that does not precede h and such that $s_j[h] \in S_j(h')$. Hence,

$$s_{-i}[h] \in S_{-i}^k(h') \text{ for every } h' \text{ that does not precede } h \tag{B.7}$$

and such that $s_{-i}[h] \in S_{-i}(h')$.

Since $s_{-i}[h] \in S_{-i}(h)$, it follows in particular that $s_{-i}[h] \in S_{-i}^k(h)$.

Let $h_i^0, h_i^1, \dots, h_i^M$ be a numbering of the information sets of player i which respects their precedence ordering. That is, if h_i^l precedes h_i^m then $l < m$. Hence, it must be that $h_i^0 = \emptyset$. Let b_i^{opp} be the cautious non-standard belief about the opponents' strategy combinations given by

$$b_i^{opp}(s_{-i}) = \begin{cases} 1 - a, & \text{if } s_{-i} = s_{-i}[h_i^0] \\ \varepsilon^m, & \text{if } s_{-i} \neq s_{-i}[h_i^0], \text{ and } m \in \{1, \dots, M\} \text{ is minimal with } s_{-i} = s_{-i}[h_i^m], \\ \varepsilon^{M+1}, & \text{otherwise,} \end{cases}$$

where a is chosen such that $\sum_{s_{-i} \in S_{-i}} b_i^{opp}(s_{-i}) = 1$. Hence, $st(a) = 0$.

We will now show that $b_i^{opp}(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$. Take some arbitrary $h \in H_i$, and let $h = h_i^m$. Let $l \in \{0, 1, \dots, M\}$ be the smallest number such that $s_{-i}[h_i^l] \in S_{-i}(h_i^m)$. Then, by construction, the standard part of the conditional belief $b_i^{opp}(h_i^m)$ assigns probability 1 to $s_{-i}[h_i^l]$. That is, $b_i^{opp}(h_i^m)$ believes $\{s_{-i}[h_i^l]\}$. Since $s_{-i}[h_i^l] \in S_{-i}(h_i^m)$, we know that $l \leq m$, and hence h_i^m does not precede h_i^l . We thus conclude that $s_{-i}[h_i^l] \in S_{-i}(h_i^m)$ and that h_i^m does not precede h_i^l . But then, by (B.7), $s_{-i}[h_i^l] \in S_{-i}^k(h_i^m)$. As the conditional belief $b_i^{opp}(h_i^m)$ believes $\{s_{-i}[h_i^l]\}$, we conclude that $b_i^{opp}(h_i^m)$ believes $S_{-i}^k(h_i^m)$. This holds for every m , and hence $b_i^{opp}(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$.

In this way, we can construct a cautious non-standard belief b_i^{opp} on the opponents' strategy combinations such that $b_i^{opp}(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$. With b_i^{self} at hand, we can then define the belief b_i^{self} in the same way as above, guaranteeing that

$b_i = (b_i^{self}, b_i^{opp})$ believes in his own future rationality and deems his own mistakes least likely. Hence, $b_i \in B_i^0$. Since, moreover, $b_i^{opp}(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$, we conclude that $b_i \in B_i^k$. We have thus shown that B_i^k is non-empty.

By induction on k , it follows that $S_i^k(h)$ and B_i^k are always non-empty for every player i , every information set h and every $k \in \mathbb{N}_0$. In particular, $S_i^k(\emptyset)$ is always non-empty for all $k \in \mathbb{N}_0$. Since the procedure terminates within finitely many steps, it follows that for every player i there is at least one perfect backwards rationalizable strategy. \square

Appendix C. Proof of Theorem 4.1

To prove Theorem 4.1, we proceed by three preparatory steps.

For the first step, consider a belief pair (b_i^{self}, b_i^{opp}) in B_i . For every information set $h \in H_i$ and choice $c_i \in C_i(h)$, let $st(b_i^{self}(h, c_i))$ be the standard part of the conditional belief $b_i^{self}(h, c_i)$ on $S_i(h, c_i)$. Moreover, let $tr(b_i^{opp})$ be the minimal cautious truncation of the cautious belief b_i^{opp} on S_{-i} , as defined in Section A.3. For every $h \in H_i$, this truncated belief $tr(b_i^{opp})$ induces a conditional cautious belief $tr(b_i^{opp})(h)$ on $S_{-i}(h)$. We say that a choice $c_i^* \in C_i(h)$ is *locally rational* for $((st(b_i^{self}(h, c_i)))_{c_i \in C_i(h)}, tr(b_i^{opp}))$ at h if

$$\begin{aligned} & \sum_{s_i \in S_i(h, c_i^*)} \sum_{s_{-i} \in S_{-i}(h)} st(b_i^{self}(h, c_i^*))(s_i) \cdot tr(b_i^{opp})(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ & \geq \sum_{s_i \in S_i(h, c_i)} \sum_{s_{-i} \in S_{-i}(h)} st(b_i^{self}(h, c_i))(s_i) \cdot tr(b_i^{opp})(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \end{aligned}$$

for every $c_i \in C_i(h)$.

Lemma C.1 (Truncation preserves local rationality). *Let (b_i^{self}, b_i^{opp}) be a belief pair in B_i that deems own mistakes least likely, $h \in H_i$ an information set for player i , and $c_i^* \in C_i(h)$ a choice for player i at h . If c_i^* is locally rational for (b_i^{self}, b_i^{opp}) at h , then c_i^* is also locally rational at h for the truncated beliefs $((st(b_i^{self}(h, c_i)))_{c_i \in C_i(h)}, tr(b_i^{opp}))$.*

Proof. Let the power series representation of the belief b_i^{opp} on S_{-i} , as defined in Section A.3, be given by

$$b_i^{opp} = p_0 + \epsilon p_1 + \epsilon^2 p_2 + \dots,$$

and let the minimal cautious truncation on S_{-i} be

$$tr(b_i^{opp}) = p_0 + \epsilon p_1 + \dots + \epsilon^K p_K.$$

By definition of the minimal cautious truncation, there must be some $s_{-i}^* \in S_{-i}$ such that

$$b_i^{opp}(s_{-i}^*) = \epsilon^K p_K(s_{-i}^*) + \epsilon^{K+1} p_{K+1}(s_{-i}^*) + \dots \tag{C.1}$$

with $p_K(s_{-i}^*) > 0$.

Then, the conditional beliefs at h induced by b_i^{opp} and $tr(b_i^{opp})$ are given by

$$b_i^{opp}(h)(s_{-i}) = \frac{1}{a}(p_0(s_{-i}) + \epsilon p_1(s_{-i}) + \epsilon^2 p_2(s_{-i}) + \dots) \tag{C.2}$$

for every $s_{-i} \in S_{-i}(h)$, where $a := \sum_{s_{-i} \in S_{-i}(h)} b_i^{opp}(s_{-i})$, and

$$tr(b_i^{opp})(h)(s_{-i}) = \frac{1}{b}(p_0(s_{-i}) + \epsilon p_1(s_{-i}) + \dots + \epsilon^K p_K(s_{-i})) \tag{C.3}$$

for every $s_{-i} \in S_{-i}(h)$, where $b := \sum_{s_{-i} \in S_{-i}(h)} tr(b_i^{opp})(s_{-i})$.

For every choice $c_i \in C_i(h)$, let the power series representation of the conditional belief $b_i^{self}(h, c_i)$ on $S_i(h, c_i)$ be given by

$$b_i^{self}(h, c_i) = q_0^{c_i} + \epsilon q_1^{c_i} + \epsilon^2 q_2^{c_i} + \dots,$$

which implies that

$$b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i))(s_i) = \epsilon q_1^{c_i}(s_i) + \epsilon^2 q_2^{c_i}(s_i) + \dots \tag{C.4}$$

for every $s_i \in S_i(h, c_i)$.

As (b_i^{self}, b_i^{opp}) deems own mistakes least likely, we must have that

$b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i))(s_i)$ is of infinitely smaller size than $b_i^{opp}(s_{-i}^*)$ for every $c_i \in C_i(h)$ and every $s_i \in S_i(h, c_i)$. By (C.1) and (C.4) it thus follows that

$$b_i^{self}(h, c_i)(s_i) - st(b_i^{self}(h, c_i))(s_i) = \epsilon^{K+1} q_{K+1}^{c_i}(s_i) + \epsilon^{K+2} q_{K+2}^{c_i}(s_i) + \dots,$$

and hence

$$b_i^{self}(h, c_i) = q_0^{c_i} + \varepsilon^{K+1} q_{K+1}^{c_i} + \varepsilon^{K+2} q_{K+2}^{c_i} + \dots \tag{C.5}$$

Let $b_i = (b_i^{self}, b_i^{opp})$. By (C.2) and (C.5) it follows, for every choice $c_i \in C_i(h)$, that

$$\begin{aligned} u_i(c_i, b_i, h) &= \sum_{s_i \in S_i(h, c_i)} \sum_{s_{-i} \in S_{-i}(h)} b_i^{self}(h, c_i)(s_i) \cdot b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ &= \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} b_i^{self}(h, c_i)(s_i) \cdot b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ &= \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left(q_0^{c_i}(s_i) + \varepsilon^{K+1} q_{K+1}^{c_i}(s_i) + \varepsilon^{K+2} q_{K+2}^{c_i}(s_i) + \dots \right) \cdot \\ &\quad \cdot \left(p_0(s_{-i}) + \varepsilon p_1(s_{-i}) + \varepsilon^2 p_2(s_{-i}) + \dots \right) \cdot u_i(z(s_i, s_{-i})) \\ &= \frac{1}{a} \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left(q_0^{c_i}(s_i) p_0(s_{-i}) + \varepsilon q_0^{c_i}(s_i) p_1(s_{-i}) + \dots \right. \\ &\quad \left. + \varepsilon^K q_0^{c_i}(s_i) p_K(s_{-i}) + d^{c_i}(s_i, s_{-i}) \right) \cdot u_i(z(s_i, s_{-i})), \end{aligned}$$

where $v(d^{c_i}(s_i, s_{-i})) \geq K + 1$. Remember that $v(d^{c_i}(s_i, s_{-i}))$ denotes the valuation of $d^{c_i}(s_i, s_{-i})$, which is the index of the leading coefficient in the power series representation of $d^{c_i}(s_i, s_{-i})$. Moreover, recall that the choice c_i^* is locally rational for $b_i = (b_i^{self}, b_i^{opp})$ at h . Then, for every choice $c_i \in C_i(h)$,

$$\begin{aligned} u_i(c_i^*, b_i, h) - u_i(c_i, b_i, h) &= \frac{1}{a} \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left((q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_0(s_{-i}) + \right. \\ &\quad \left. + \varepsilon (q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_1(s_{-i}) + \dots \right. \\ &\quad \left. + \varepsilon^K (q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_K(s_{-i}) + \hat{d}^{c_i}(s_i, s_{-i}) \right) \cdot u_i(z(s_i, s_{-i})), \end{aligned} \tag{C.6}$$

where $v(\hat{d}^{c_i}(s_i, s_{-i})) \geq K + 1$.

Since c_i^* is locally rational for $b_i = (b_i^{self}, b_i^{opp})$ at h , we have that $u_i(c_i^*, b_i, h) - u_i(c_i, b_i, h) \geq 0$ for all $c_i \in C_i(h)$. This implies that $\frac{a}{b} \cdot (u_i(c_i^*, b_i, h) - u_i(c_i, b_i, h)) \geq 0$. By Lemma A.1 we thus know that $trunc^K(\frac{a}{b} \cdot (u_i(c_i^*, b_i, h) - u_i(c_i, b_i, h))) \geq 0$.

By (C.6) we thus conclude that

$$\begin{aligned} trunc^K\left(\frac{a}{b} \cdot (u_i(c_i^*, b_i, h) - u_i(c_i, b_i, h))\right) &= \frac{1}{b} \left(\sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left((q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_0(s_{-i}) + \right. \right. \\ &\quad \left. \left. + \varepsilon (q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_1(s_{-i}) + \dots + \varepsilon^K (q_0^{c_i^*}(s_i) - q_0^{c_i}(s_i)) p_K(s_{-i}) \right) \cdot u_i(z(s_i, s_{-i})) \right) \geq 0 \end{aligned}$$

for every $c_i \in C_i(h)$. Hence,

$$\begin{aligned} &\frac{1}{b} \cdot \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left(q_0^{c_i^*}(s_i) p_0(s_{-i}) + \varepsilon q_0^{c_i^*}(s_i) p_1(s_{-i}) + \dots + \varepsilon^K q_0^{c_i^*}(s_i) p_K(s_{-i}) \right) \cdot u_i(z(s_i, s_{-i})) \\ &\geq \frac{1}{b} \cdot \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} \left(q_0^{c_i}(s_i) p_0(s_{-i}) + \varepsilon q_0^{c_i}(s_i) p_1(s_{-i}) + \dots + \varepsilon^K q_0^{c_i}(s_i) p_K(s_{-i}) \right) \cdot u_i(z(s_i, s_{-i})) \end{aligned}$$

for all $c_i \in C_i(h)$. As $st(b_i^{self}(h, c_i)) = q_0^{c_i}$ for all $c_i \in C_i(h)$ and $tr(b_i^{opp})(h)$ is given by (C.3), the above inequality is equivalent to

$$\begin{aligned} &\sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} st(b_i^{self}(h, c_i^*))(s_i) \cdot tr(b_i^{opp})(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ &\geq \sum_{s_i \in S_i(h)} \sum_{s_{-i} \in S_{-i}(h)} st(b_i^{self}(h, c_i))(s_i) \cdot tr(b_i^{opp})(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \end{aligned}$$

for all $c_i \in C_i(h)$. This, in turn, means that c_i^* is locally rational at h for $((st(b_i^{self}(h, c_i)))_{c_i \in C_i(h)}, tr(b_i^{opp}))$, which was to show. \square

As a second step, we prove that if a player believes in his own future rationality, then local rationality of a strategy at all information sets weakly following information set h^* implies global rationality of this strategy at h^* . To define this lemma formally, we need some additional notation and definitions. Let $\hat{b}_i^{self} = (\hat{b}_i^{self}(h, c_i))_{h \in H_i, c_i \in C_i(h)}$, where $\hat{b}_i^{self}(h, c_i)$ is a standard probability distribution on $S_i(h, c_i)$ for every $h \in H_i$ and every $c_i \in C_i(h)$. Moreover, let $b_i^{opp} \in B_i^{opp}$. For a given information set $h \in H_i$ and choice $c_i \in C_i(h)$, we define the expected utility

$$u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h) := \sum_{s_i \in S_i(h, c_i)} \sum_{s_{-i} \in S_{-i}(h)} \hat{b}_i^{self}(h, c_i)(s_i) \cdot b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})).$$

We call c_i locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at h if

$$u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h) \geq u_i(c'_i, (\hat{b}_i^{self}, b_i^{opp}), h) \text{ for all } c'_i \in C_i(h).$$

We say that $(\hat{b}_i^{self}, b_i^{opp})$ believes in his own future rationality if for every $h \in H_i$ and $c_i \in C_i(h)$, the standard probability distribution $\hat{b}_i^{self}(h, c_i)$ only assigns positive probability to strategies $s_i \in S_i(h, c_i)$ where $s_i(h')$ is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at every $h' \in H_i(s_i)$ following h .

Lemma C.2 (When local rationality implies global rationality). Let

$\hat{b}_i^{self} = (\hat{b}_i^{self}(h, c_i))_{h \in H_i, c_i \in C_i(h)}$ where $\hat{b}_i^{self}(h, c_i)$ is a standard probability distribution on $S_i(h, c_i)$ for every $h \in H_i$ and every $c_i \in C_i(h)$. Let $b_i^{opp} \in B_i^{opp}$ and assume that $(\hat{b}_i^{self}, b_i^{opp})$ believes in his own future rationality. Let $s_i^* \in S_i$ and $h^* \in H_i(s_i^*)$ such that $s_i^*(h)$ is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at every $h \in H_i(s_i^*)$ weakly following h^* . Then, s_i^* is globally rational for b_i^{opp} at h^* .

Proof. We first introduce some additional notation. For every information set $h \in H_i$ and every choice $c_i \in C_i(h)$, let

$$u_i^{\max}(b_i^{opp}, h) := \max_{s_i \in S_i(h)} u_i(s_i, b_i^{opp}(h))$$

and

$$u_i^{\max}(b_i^{opp}, h, c_i) := \max_{s_i \in S_i(h, c_i)} u_i(s_i, b_i^{opp}(h)).$$

Then, we have that

$$u_i^{\max}(b_i^{opp}, h) = \max_{c_i \in C_i(h)} u_i^{\max}(b_i^{opp}, h, c_i) \tag{C.7}$$

for every $h \in H_i$. Moreover, strategy s_i is globally rational for b_i^{opp} at $h \in H_i(s_i)$ if

$$u_i(s_i, b_i^{opp}(h)) = u_i^{\max}(b_i^{opp}, h).$$

For every information set $h \in H_i$ and every choice $c_i \in C_i(h)$ we also define

$$u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h) := \sum_{s_i \in S_i(h, c_i)} \sum_{s_{-i} \in S_{-i}(h)} \hat{b}_i^{self}(h, c_i)(s_i) \cdot b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})).$$

Hence, strategy s_i is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at $h \in H_i(s_i)$ if

$$u_i(s_i(h), (\hat{b}_i^{self}, b_i^{opp}), h) \geq u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h) \text{ for all } c_i \in C_i(h).$$

We prove the statement in the lemma by induction on the number of player i information sets that follow h^* . If h^* is not followed by any player i information set, then the statement holds because local rationality for $(\hat{b}_i^{self}, b_i^{opp})$ at h^* coincides with global rational for b_i^{opp} at h .

Suppose now that h^* is followed by $k \in \mathbb{N}$ consecutive player i information sets, and that the statement holds for every player i information set that follows h^* . Consider some choice $c_i \in C_i(h^*)$. By $H_i^+(h^*, c_i)$ we denote the collection of information sets $h \in H_i$ such that h weakly follows h^* and c_i , and there is no $h' \in H_i$ preceding h that also weakly follows h^* and c_i . Let $S_{-i}^{not}(h^*, c_i)$ be the collection of those opponents' strategy combinations $s_{-i} \in S_{-i}(h^*)$ that after h^* and c_i do not lead to any player i information set.

Then, for every $c_i \in C_i(h^*)$ we have that

$$\begin{aligned} u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h^*) &= \sum_{s_i \in S_i(h^*, c_i)} \sum_{s_{-i} \in S_{-i}(h^*)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ &= \sum_{s_i \in S_i(h^*, c_i)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot \\ &\quad \cdot \left[\sum_{h \in H_i^+(h^*, c_i)} \sum_{s_{-i} \in S_{-i}(h)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) + \right. \\ &\quad \left. + \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right] \\ &= \sum_{s_i \in S_i(h^*, c_i)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot \\ &\quad \cdot \left[\sum_{h \in H_i^+(h^*, c_i)} b_i^{opp}(h^*)(S_{-i}(h)) \sum_{s_{-i} \in S_{-i}(h)} \frac{b_i^{opp}(h^*)(s_{-i})}{b_i^{opp}(h^*)(S_{-i}(h))} \cdot u_i(z(s_i, s_{-i})) \right] \end{aligned}$$

$$\begin{aligned}
 & + \left. \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right] \\
 = & \sum_{s_i \in S_i(h^*, c_i)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot \\
 & \left[\sum_{h \in H_i^+(h^*, c_i)} b_i^{opp}(h^*)(S_{-i}(h)) \sum_{s_{-i} \in S_{-i}(h)} b_i^{opp}(h)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right. \\
 & \left. + \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right] \\
 = & \sum_{s_i \in S_i(h^*, c_i)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot \left[\sum_{h \in H_i^+(h^*, c_i)} b_i^{opp}(h^*)(S_{-i}(h)) \cdot u_i(s_i, b_i^{opp}(h)) \right. \\
 & \left. + \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right]. \tag{C.8}
 \end{aligned}$$

Here, the fourth equality follows from the rules of conditional probabilities.

As $(\hat{b}_i^{self}, b_i^{opp})$ believes in his own future rationality, $\hat{b}_i^{self}(h^*, c_i)$ only assigns positive probability to $s_i \in S_i(h^*, c_i)$ where s_i is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at every $h \in H_i^+(h^*, c_i)$, and every $h' \in H_i(s_i)$ that follows h . By the induction assumption, we know that every such s_i is globally rational at every $h \in H_i^+(h^*, c_i)$. Hence, $\hat{b}_i^{self}(h^*, c_i)$ only assigns positive probability to $s_i \in S_i(h^*, c_i)$ where

$$u_i(s_i, b_i^{opp}(h)) = u_i^{\max}(b_i^{opp}, h)$$

for every $h \in H_i^+(h^*, c_i)$. Together with (C.8) we conclude that

$$\begin{aligned}
 u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h^*) & = \sum_{s_i \in S_i(h^*, c_i)} \hat{b}_i^{self}(h^*, c_i)(s_i) \cdot \left[\sum_{h \in H_i^+(h^*, c_i)} b_i^{opp}(h^*)(S_{-i}(h)) \cdot u_i^{\max}(b_i^{opp}, h) \right. \\
 & \left. + \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \right] \\
 & = \sum_{h \in H_i^+(h^*, c_i)} b_i^{opp}(h^*)(S_{-i}(h)) \cdot u_i^{\max}(b_i^{opp}, h) + \\
 & \quad + \sum_{s_{-i} \in S_{-i}^{not}(h^*, c_i)} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \tag{C.9} \\
 & = u_i^{\max}(b_i^{opp}, h^*, c_i).
 \end{aligned}$$

Here, the last equality follows from the fact that the terminal node $z(s_i, s_{-i})$ does not depend on the specific $s_i \in S_i(h^*, c_i)$ if $s_{-i} \in S_{-i}^{not}(h^*, c_i)$. Hence, we see that

$$u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h^*) = u_i^{\max}(b_i^{opp}, h^*, c_i) \text{ for all } c_i \in C_i(h^*). \tag{C.10}$$

As $s_i^*(h^*)$ is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at h^* , we know that

$$\begin{aligned}
 u_i(s_i^*(h^*), (\hat{b}_i^{self}, b_i^{opp}), h^*) & = \max_{c_i \in C_i(h^*)} u_i(c_i, (\hat{b}_i^{self}, b_i^{opp}), h^*) \\
 & = \max_{c_i \in C_i(h^*)} u_i^{\max}(b_i^{opp}, h^*, c_i) = u_i^{\max}(b_i^{opp}, h^*), \tag{C.11}
 \end{aligned}$$

where the second equality follows from (C.10) and the last equality from (C.7).

On the other hand, we know by (C.9) that

$$\begin{aligned}
 u_i(s_i^*(h^*), (\hat{b}_i^{self}, b_i^{opp}), h^*) & = \sum_{h \in H_i^+(h^*, s_i^*(h^*))} b_i^{opp}(h^*)(S_{-i}(h)) \cdot u_i^{\max}(b_i^{opp}, h) + \\
 & \quad + \sum_{s_{-i} \in S_{-i}^{not}(h^*, s_i^*(h^*))} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})). \tag{C.12}
 \end{aligned}$$

As $s_i^*(h)$ is assumed to be locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at every $h \in H_i(s_i^*)$ weakly following h^* , we know that, for every $h \in H_i^+(h^*, s_i^*(h^*))$, the choice $s_i^*(h')$ is locally rational for $(\hat{b}_i^{self}, b_i^{opp})$ at every $h' \in H_i(s_i^*)$ weakly following h . Hence, by the induction assumption, s_i^* is globally rational for b_i^{opp} at every $h \in H_i^+(h^*, s_i^*(h^*))$, which means that

$$u_i(s_i^*, b_i^{opp}(h)) = u_i^{\max}(b_i^{opp}, h) \text{ for every } h \in H_i^+(h^*, s_i^*(h^*)). \tag{C.13}$$

By combining (C.12) and (C.13) we obtain that

$$\begin{aligned} u_i(s_i^*(h^*), (\hat{b}_i^{self}, b_i^{opp}), h^*) &= \sum_{h \in H_i^+(h^*, s_i^*(h^*))} b_i^{opp}(h^*)(S_{-i}(h)) \cdot u_i(s_i^*, b_i^{opp}(h)) + \\ &\quad + \sum_{s_{-i} \in S_{-i}^{not}(h^*, s_i^*(h^*))} b_i^{opp}(h^*)(s_{-i}) \cdot u_i(z(s_i, s_{-i})) \\ &= u_i(s_i^*, b_i^{opp}(h^*)). \end{aligned} \tag{C.14}$$

From (C.14) and (C.11) we can thus conclude that

$$u_i(s_i^*, b_i^{opp}(h^*)) = u_i(s_i^*(h^*), (\hat{b}_i^{self}, b_i^{opp}), h^*) = u_i^{\max}(b_i^{opp}, h^*).$$

This means that s_i^* is globally rational for b_i^{opp} at h^* , which was to show. By induction, the proof is thus complete. \square

As a third step, we are able to derive the following important result by combining Lemma C.1 and Lemma C.2. This step will be crucial for proving our main theorem below.

Corollary C.1 (From local to global rationality). *Let (b_i^{self}, b_i^{opp}) be a belief pair in B_i that deems own mistakes least likely and believes in his own future rationality. Let $s_i^* \in S_i$ and $h^* \in H_i(s_i^*)$ such that s_i^* is locally rational for (b_i^{self}, b_i^{opp}) at every $h \in H_i(s_i^*)$ that weakly follows h^* . Then, s_i^* is globally rational at h^* for the minimal cautious truncation $tr(b_i^{opp})$ of b_i^{opp} .*

Proof. Suppose that s_i^* is locally rational for (b_i^{self}, b_i^{opp}) at every $h \in H_i(s_i^*)$ that weakly follows h^* . That is, $s_i^*(h)$ is locally rational for (b_i^{self}, b_i^{opp}) at every $h \in H_i(s_i^*)$ that weakly follows h^* . Since (b_i^{self}, b_i^{opp}) deems own mistakes least likely, it follows from Lemma C.1 that $s_i^*(h)$ is also locally rational for the truncated belief pair $((st(b_i^{self}(h, c_i)))_{c_i \in C_i(h)}, tr(b_i^{opp}))$ at every $h \in H_i(s_i^*)$ that weakly follows h^* .

Define $\hat{b}_i^{self} := (st(b_i^{self}(h, c_i)))_{h \in H_i, c_i \in C_i(h)}$. We will show that $(\hat{b}_i^{self}, tr(b_i^{opp}))$ believes in his own future rationality. As the original belief pair (b_i^{self}, b_i^{opp}) believes in his own future rationality, we know that for every $h \in H_i$ and $c_i \in C_i(h)$, the standard part of $b_i^{self}(h, c_i)$ only assigns positive probability to strategies $s_i \in S_i(h, c_i)$ where $s_i(h')$ is locally rational for (b_i^{self}, b_i^{opp}) at every $h' \in H_i(s_i)$ following h . By Lemma C.1 we know that every such $s_i(h')$ is also locally rational for $(\hat{b}_i^{self}, tr(b_i^{opp}))$ at h' . Hence, we conclude that for every $h \in H_i$ and $c_i \in C_i(h)$, the belief $\hat{b}_i^{self}(h, c_i) = st((b_i^{self}(h, c_i)))$ only assigns positive probability to strategies $s_i \in S_i(h, c_i)$ where $s_i(h')$ is locally rational for $(\hat{b}_i^{self}, tr(b_i^{opp}))$ at every $h' \in H_i(s_i)$ following h . Therefore, $(\hat{b}_i^{self}, tr(b_i^{opp}))$ believes in his own future rationality.

As such, we conclude that $s_i^*(h)$ is locally rational for the truncated belief pair $(\hat{b}_i^{self}, tr(b_i^{opp}))$ at every $h \in H_i(s_i^*)$ that weakly follows h^* , and that $(\hat{b}_i^{self}, tr(b_i^{opp}))$ believes in his own future rationality. By Lemma C.2 it follows that s_i^* is globally rational for $tr(b_i^{opp})$ at h^* , which was to show. \square

We are now fully equipped to prove Theorem 4.1.

Proof of Theorem 4.1. For every $k \in \mathbb{N}_0$, every player i and every $h \in H_i$, let $S_{i,qp}^{k}(h)$ and $B_{i,qp}^{opp,k}$ be the sets of strategies and beliefs that survive round k of the procedural quasi-perfect rationalizability procedure. Similarly, let $S_{i,ss}^k(h)$ and $B_{i,ss}^k$ be the sets of strategies and belief pairs that survive round k of the perfect backwards rationalizability procedure. As before, for every $b_i^{opp} \in B_i^{opp}$ we denote by $tr(b_i^{opp})$ the minimal cautious truncation of b_i^{opp} on S_{-i} . We prove the following claim.

Claim. *For every $k \in \mathbb{N}_0$, every player i and every $h^* \in H_i$, (a) $S_{i,ss}^k(h^*) \subseteq S_{i,qp}^k(h^*)$, and (b) for every $(b_i^{self}, b_i^{opp}) \in B_{i,ss}^k$ it holds that $tr(b_i^{opp}) \in B_{i,qp}^{opp,k}$.*

Proof of claim. We prove so by induction on k . For $k = 0$ the statement is trivial since $S_{i,ss}^0(h^*) = S_{i,qp}^0(h^*) = S_i(h^*)$ and $B_{i,qp}^{opp,0}$ is the set of all cautious non-standard probability distributions on S_{-i} .

Let $k \in \mathbb{N}$, and suppose that (a) and (b) are true for $k - 1$. To show (a) for k , take some strategy $s_i^* \in S_{i,ss}^k(h^*)$. Then, by definition, $s_i^* \in S_{i,ss}^{k-1}(h^*)$, and there is some $(b_i^{self}, b_i^{opp}) \in B_{i,ss}^{k-1}$ such that $s_i^*(h)$ is locally rational for (b_i^{self}, b_i^{opp}) at every $h \in H_i(s_i^*)$ weakly following h^* . Since, by the induction assumption on (a), $S_{i,ss}^{k-1}(h^*) \subseteq S_{i,qp}^{k-1}(h^*)$, we know that $s_i^* \in S_{i,qp}^{k-1}(h^*)$. Moreover, as $(b_i^{self}, b_i^{opp}) \in$

$B_{i,ss}^{k-1} \subseteq B_{i,ss}^0$ we know that (b_i^{self}, b_i^{opp}) deems own mistakes least likely and believes in his own future rationality. Since s_i^* is locally rational for (b_i^{self}, b_i^{opp}) at every $h \in H_i(s_i^*)$ weakly following h^* , we thus conclude by Corollary C.1 that s_i^* is globally rational for $tr(b_i^{opp})$ at every $h \in H_i(s_i^*)$ weakly following h^* . Moreover, as $(b_i^{self}, b_i^{opp}) \in B_{i,ss}^{k-1}$, we know by the induction assumption on (b) that $tr(b_i^{opp}) \in B_{i,qp}^{opp,k-1}$.

Summarizing, we see that $s_i^* \in S_{i,qp}^{k-1}(h^*)$, and that s_i^* is globally rational for $tr(b_i^{opp}) \in B_{i,qp}^{opp,k-1}$ at every $h \in H_i(s_i^*)$ weakly following h^* . Hence, by definition, $s_i^* \in S_{i,qp}^k(h^*)$. We thus conclude that $S_{i,ss}^k(h^*) \subseteq S_{i,qp}^k(h^*)$.

To show (b), take some $(b_i^{self}, b_i^{opp}) \in B_{i,ss}^k$. Then, by definition, $(b_i^{self}, b_i^{opp}) \in B_{i,ss}^{k-1}$ and $b_i^{opp}(h)$ believes $S_{-i,ss}^{k-1}(h)$ for every $h \in H_i$. By the induction assumption on (b) we already know that $tr(b_i^{opp}) \in B_{i,qp}^{opp,k-1}$. Since $b_i^{opp}(h)$ believes $S_{-i,ss}^{k-1}(h)$, the standard part of $b_i^{opp}(h)$ only assigns positive probability to opponents' strategy combinations $s_{-i} \in S_{-i,ss}^{k-1}(h)$. Note that the standard part of $b_i^{opp}(h)$ is the same as the standard part of $tr(b_i^{opp})(h)$. Hence, the standard part of $tr(b_i^{opp})(h)$ only assigns positive probability to $s_{-i} \in S_{-i,ss}^{k-1}(h)$. By the induction assumption on (a) we know that $S_{-i,ss}^{k-1}(h) \subseteq S_{-i,qp}^{k-1}(h)$, and therefore the standard part of $tr(b_i^{opp})(h)$ only assigns positive probability to $s_{-i} \in S_{-i,qp}^{k-1}(h)$. In other words, $tr(b_i^{opp})(h)$ believes $S_{-i,qp}^{k-1}(h)$.

Summarizing, we see that $tr(b_i^{opp}) \in B_{i,qp}^{opp,k-1}$ and that $tr(b_i^{opp})(h)$ believes $S_{-i,qp}^{k-1}(h)$ for every $h \in H_i$. Hence, by definition, $tr(b_i^{opp}) \in B_{i,qp}^{opp,k}$, as was to show.

By induction on k , (a) and (b) are true for every $k \in \mathbb{N}_0$, which completes the proof of the claim.

To prove the theorem, consider some perfect backwards rationalizable strategy s_i^* for player i . Then, by definition, $s_i^* \in S_{i,ss}^k(\emptyset)$ for all $k \in \mathbb{N}_0$. Hence, by part (a) of the claim, $s_i^* \in S_{i,qp}^k(\emptyset)$ for all $k \in \mathbb{N}_0$, which means that s_i^* is procedurally quasi-perfectly rationalizable. This completes the proof. \square

Appendix D. Relation to quasi-perfect rationalizability

In this section we will compare our notion of procedural quasi-perfect rationalizability to quasi-perfect rationalizability as defined by Asheim and Perea (2005). To that purpose, we first review the definition of quasi-perfect rationalizability, and subsequently show that in all games, every quasi-perfectly rationalizable strategy is also procedurally quasi-perfectly rationalizable in our sense. We then show by means of a counterexample that there are procedurally quasi-perfectly rationalizable strategies which are not quasi-perfectly rationalizable. Hence, quasi-perfect rationalizability is a strict refinement of procedural quasi-perfect rationalizability. The same example also demonstrates that even perfect backwards rationalizable strategy need not be quasi-perfectly rationalizable.

D.1. Quasi-perfect rationalizability

We have defined procedural quasi-perfect rationalizability by means of a procedure, that recursively eliminates strategies and beliefs from the game. Asheim and Perea (2005) take a different approach, since they define quasi-perfect rationalizability by looking at *belief hierarchies* encoded by types within an epistemic model. Also, they use *lexicographic beliefs* (Blume et al. (1991)) rather than non-standard beliefs to model cautious reasoning. That is, they take as a primitive not only beliefs about the opponents' strategies, as we do, but also beliefs about the opponents' beliefs about the other players' strategies (second-order beliefs), and higher-order beliefs. Quasi-perfect rationalizability is defined by imposing epistemic conditions on such belief hierarchies. Since we have used non-standard beliefs, rather than lexicographic beliefs, to define perfect backwards rationalizability and procedural quasi-perfect rationalizability, we will reproduce the definition of quasi-perfect rationalizability by using non-standard beliefs instead of lexicographic beliefs. This is justified, since Halpern (2010) shows in Theorem 4.5 that for every lexicographic belief on a finite set there is an "equivalent" non-standard belief on that set, and *vice versa*.

Definition D.1 (*Epistemic model with non-standard beliefs*). For a given dynamic game G , a finite epistemic model with non-standard beliefs is a tuple $M = (T_i, \beta_i)_{i \in I}$ such that, for every player i ,

- (a) T_i is a finite set of types, and
- (b) β_i is a function that assigns to every type $t_i \in T_i$ a non-standard belief $\beta_i(t_i)$ on $S_{-i} \times T_{-i}$.

In Asheim and Perea (2005), every cautious type $t_i \in T_i$ holds a system of conditional cautious lexicographic beliefs on $S_{-i} \times T_{-i}$, induced by a *single* cautious lexicographic belief $\hat{\beta}_i(t_i)$ on $S_{-i} \times T_{-i}$. That is, $\hat{\beta}_i(t_i) = (\hat{\beta}_i^1(t_i); \hat{\beta}_i^2(t_i); \dots; \hat{\beta}_i^K(t_i))$, where $\hat{\beta}_i^k(t_i) \in \Delta(S_{-i} \times T_{-i})$ for all $k \in \{1, \dots, K\}$, and where for every $t_{-i} \in T_{-i}$ that receives positive probability by $\hat{\beta}_i^k(t_i)$ for some $k \in \{1, \dots, K\}$, and every $s_{-i} \in S_{-i}$, there is some $m \in \{1, \dots, K\}$ such that $\hat{\beta}_i^m(s_{-i}, t_{-i}) > 0$. Thus, the lexicographic belief $\hat{\beta}_i(t_i)$ in Asheim and Perea (2005) plays the same role as the non-standard belief $\beta_i(t_i)$ in our setting.

An epistemic model is used to *encode* non-standard belief hierarchies for the players, including beliefs about the opponents' strategies, beliefs about the opponents' beliefs about their opponents' strategies, and so on. The concept of quasi-perfect rationalizability restricts to types that express common full belief in "caution" and the "event that types induce sequentially rational behavioral strategies". We will now formally define these events.

Definition D.2 (Caution). Consider a finite epistemic model $M = (T_i, \beta_i)_{i \in I}$ for a dynamic game G . A type $t_i \in T_i$ is cautious if for every opponents' type combination $t_{-i} \in T_{-i}$ with $\beta_i(t_i)(t_{-i}) > 0$, it holds that $\beta_i(t_i)(s_{-i}, t_{-i}) > 0$ for every $s_{-i} \in S_{-i}$.

Here, $\beta_i(t_i)(t_{-i})$ is an abbreviation for the marginal probability $\beta_i(t_i)(S_{-i} \times \{t_{-i}\})$. We will use such abbreviations for marginals more often in the remainder of this section. Hence, caution states that if t_i deems possible a type combination t_{-i} for his opponents, then he must deem possible every strategy combination for that type combination. In particular, t_i holds a cautious non-standard belief on the set S_{-i} of opponents' strategy combinations. This is similar to how Asheim and Perea (2005) define a cautious type with lexicographic beliefs, in line with our description above.

Consider a cautious type t_i and an information set $h \in H_i$. By $\beta_i(t_i, h)$ we denote the induced (cautious) conditional belief on $S_{-i}(h) \times T_{-i}$. For every strategy $s_i \in S_i(h)$ we denote by

$$u_i(s_i, t_i, h) := \sum_{(s_{-i}, t_{-i}) \in S_{-i}(h) \times T_{-i}} \beta_i(t_i, h)(s_{-i}, t_{-i}) \cdot u_i(z(s_i, s_{-i}))$$

the expected (non-standard) utility at h of choosing strategy s_i under the conditional belief $\beta_i(t_i, h)$. We say that a strategy $s_i \in S_i(h)$ is *globally rational* for the cautious type t_i at $h \in H_i$ if

$$u_i(s_i, t_i, h) \geq u_i(s'_i, t_i, h) \text{ for all } s'_i \in S_i(h).$$

To define what it means for a type to “induce a sequentially rational behavioral strategy” we need some additional terminology. A *behavioral strategy* for player i is a tuple $\sigma_i = (\sigma_i(h))_{h \in H_i}$ such that $\sigma_i(h)$ is a (standard) probability distribution on the set of choices $C_i(h)$ available for player i at h . For a behavioral strategy σ_i and a strategy $s_i \in S_i$, let

$$\sigma_i(s_i) := \prod_{h \in H_i(s_i)} \sigma_i(s_i(h)) \tag{D.1}$$

be the induced probability that σ_i assigns to the strategy s_i . For a given behavioral strategy σ_i and information set $h \in H_i$, let $\sigma_i|_h$ be the behavioral strategy that (i) at every $h' \in H_i$ preceding h assigns probability 1 to the unique choice for player i at h' leading to h , and (ii) coincides with σ_i at all other information sets. We say that a behavioral strategy σ_i is *sequentially rational* for a cautious type t_i if at every information set $h \in H_i$, we have that $\sigma_i|_h(s_i) > 0$ only if s_i is globally rational for t_i at h .

Recall that, for an information set $h \in H_i$ and a choice $c_i \in C_i(h)$, we denote by $S_i(h, c_i)$ the set of strategies $s_i \in S_i(h)$ with $s_i(h) = c_i$. For a cautious type t_i and an opponent's type t_j with $\beta_i(t_i)(t_j) > 0$, let $\sigma_j^{t_i|t_j}$ be the induced behavioral strategy for player j given by

$$\sigma_j^{t_i|t_j}(h)(c_j) := st \left(\frac{\beta_i(t_i)(S_j(h, c_j) \times \{t_j\})}{\beta_i(t_i)(S_j(h) \times \{t_j\})} \right) \tag{D.2}$$

for every information set $h \in H_j$ and every choice $c_j \in C_j(h)$.

Definition D.3 (Inducing sequentially rational behavioral strategies). Consider a finite epistemic model $M = (T_i, \beta_i)_{i \in I}$ for a dynamic game G . A cautious type $t_i \in T_i$ induces sequentially rational behavioral strategies if for every opponent $j \neq i$ and every type $t_j \in T_j$ with $\beta_i(t_i)(t_j) > 0$, the induced behavioral strategy $\sigma_j^{t_i|t_j}$ is sequentially rational for t_j .

We are now ready to define quasi-perfectly rationalizable types as those types that are cautious, induce sequentially rational behavioral strategies, and express common full belief in these two events. Formally, a type t_i expresses *1-fold full belief* in caution and the event that types induce sequentially rational behavioral strategies if $\beta_i(t_i)$ only assigns positive (non-standard) probability to opponents' types that are cautious and induce sequentially rational behavioral strategies. For every $k \geq 2$, type t_i expresses *k-fold full belief* in caution and the event that types induce sequentially rational behavioral strategies if $\beta_i(t_i)$ only assigns positive (non-standard) probability to opponents' types that express $(k - 1)$ -fold full belief in caution and the event that types induce sequentially rational behavioral strategies. A type t_i expresses *common full belief* in caution and the event that types induce sequentially rational behavioral strategies if t_i expresses k -fold full belief in caution and the event that types induce sequentially rational behavioral strategies, for every $k \in \mathbb{N}$.

Definition D.4 (Quasi-perfect rationalizability). Consider a finite epistemic model $M = (T_i, \beta_i)_{i \in I}$ for a dynamic game G . A type $t_i \in T_i$ is quasi-perfectly rationalizable if it is cautious, induces sequentially rational behavioral strategies, and expresses common full belief in caution and the event that types induce sequentially rational behavioral strategies. A strategy $s_i \in S_i$ is quasi-perfectly rationalizable if there is a finite epistemic model $M = (T_i, \beta_i)_{i \in I}$ and a quasi-perfectly rationalizable type $t_i \in T_i$, such that s_i is globally rational for t_i at every $h \in H_i(s_i)$.

In the following subsection we will show that every quasi-perfectly rationalizable strategy is procedurally quasi-perfectly rationalizable in our sense, but not *vice versa*.

D.2. Relation between the two quasi-perfect rationalizability concepts

We first show that, in all dynamic games, every strategy that is quasi-perfectly rationalizable is also procedurally quasi-perfectly rationalizable in our sense.

Theorem D.1 (Relation with quasi-perfect rationalizability). *Consider a dynamic game G . Then, every strategy that is quasi-perfectly rationalizable is also procedurally quasi-perfectly rationalizable.*

Proof. Let $S_i^k(h)$ and $B_i^{opp,k}$ be the sets of strategies and beliefs that survive round k of the procedural quasi-perfect rationalizability procedure. Consider a finite epistemic model $M = (T_i, \beta_i)_{i \in I}$ for G , as in Asheim and Perea (2005). We prove, by induction on k , that for every player i , every quasi-perfectly rationalizable type $t_i \in T_i$, and every information set $h \in H_i$, we have that (a) every strategy $s_i \in S_i(h)$ that is globally rational for t_i at every $h' \in H_i(s_i)$ weakly following h is in $S_i^k(h)$, and (b) the marginal of $\beta_i(t_i)$ on S_{-i} is in $B_i^{opp,k}$.

For $k = 0$ this statement is true because $S_i^0(h) = S_i(h)$, the type t_i is cautious, and $B_i^{opp,0} = B_i^{opp}$ contains all cautious beliefs on S_{-i} .

Now let $k \in \mathbb{N}$ and suppose that (a) and (b) are true for $k - 1$ and all players i . Consider a player i , a quasi-perfectly rationalizable type $t_i \in T_i$, and an information set $h \in H_i$. To show (a), take some strategy $s_i \in S_i(h)$ that is globally rational for t_i at every $h' \in H_i(s_i)$ weakly following h . By the induction assumption on (a) we know that $s_i \in S_i^{k-1}(h)$. Let $b_i^{opp}(t_i)$ be the marginal of $\beta_i(t_i)$ on S_{-i} . By the induction assumption on (b) we know that $b_i^{opp}(t_i) \in B_i^{opp,k-1}$. Hence, $s_i \in S_i^{k-1}(h)$ is globally rational for $b_i^{opp}(t_i) \in B_i^{opp,k-1}$ at every $h' \in H_i(s_i)$ weakly following h . This implies that $s_i \in S_i^k(h)$, which completes the induction step for (a).

To show (b), let $b_i^{opp}(t_i)$ be the marginal of $\beta_i(t_i)$ on S_{-i} . By the induction assumption on (b) we know that $b_i^{opp}(t_i) \in B_i^{opp,k-1}$. To show that $b_i^{opp}(t_i) \in B_i^{opp,k}$, we must show that $b_i^{opp}(t_i)(h)$ believes $S_{-i}^k(h)$ for all $h \in H_i$. That is, we must show that $st(b_i^{opp}(t_i)(h)(s_{-i})) > 0$ only if $s_{-i} \in S_{-i}^k(h)$.

Consider some information set $h \in H_i$ and some opponents' strategy combination s_{-i} such that $st(b_i^{opp}(t_i)(h)(s_{-i})) > 0$. We will show that $s_{-i} \in S_{-i}^k(h)$. As, by the induction assumption on (b), $b_i^{opp}(t_i) \in B_i^{opp,k-1}$, it follows that $b_i^{opp}(t_i)(h)$ believes $S_{-i}^{k-1}(h)$, and hence $s_{-i} \in S_{-i}^{k-1}(h)$. Let $s_{-i} = (s_j)_{j \neq i}$. To show that $s_{-i} \in S_{-i}^k(h)$, we will show that for every opponent $j \neq i$ there is some $b_j^{opp} \in B_j^{opp,k-1}$ such that s_j is globally rational for b_j^{opp} at every $h' \in H_j(s_j)$ weakly following h .

Fix an opponent j . Since $st(b_i^{opp}(t_i)(h)(s_j)) > 0$, the belief $b_i^{opp}(t_i)$ is the marginal of $\beta_i(t_i)$ on S_{-i} , and $b_i^{opp}(t_i)(h)$ is the induced conditional belief on $S_{-i}(h)$, there must be some type $t_j \in T_j$ with $\beta_i(t_i)(t_j) > 0$ such that

$$st \left(\frac{\beta_i(t_i)(s_j, t_j)}{\beta_i(t_i)(S_j(h) \times \{t_j\})} \right) > 0. \tag{D.3}$$

Now, let $b_j^{opp}(t_j)$ be the marginal of $\beta_j(t_j)$ on S_{-j} . We show that $b_j^{opp}(t_j) \in B_j^{opp,k-1}$ and that s_j is globally rational for $b_j^{opp}(t_j)$ at every $h' \in H_j(s_j)$ weakly following h .

As $\beta_i(t_i)(t_j) > 0$ and t_i is quasi-perfectly rationalizable, it must be that t_j is quasi-perfectly rationalizable as well. Hence, by our induction assumption on (b) we conclude that $b_j^{opp}(t_j) \in B_j^{opp,k-1}$.

Consider now some $h' \in H_j(s_j)$ weakly following h . We show that s_j is globally rational for $b_j^{opp}(t_j)$ at h' . Take some arbitrary $h'' \in H_j(s_j)$ weakly following h' . Then, h'' weakly follows h and hence $S_j(h'') \subseteq S_j(h)$. Moreover, $s_j \in S_j(h''), s_j(h'')$. It thus follows by (D.3) that

$$\begin{aligned} st \left(\frac{\beta_i(t_i)(S_j(h''), s_j(h'')) \times \{t_j\}}{\beta_i(t_i)(S_j(h'') \times \{t_j\})} \right) &\geq st \left(\frac{\beta_i(t_i)(S_j(h''), s_j(h'')) \times \{t_j\}}{\beta_i(t_i)(S_j(h) \times \{t_j\})} \right) \\ &\geq st \left(\frac{\beta_i(t_i)(s_j, t_j)}{\beta_i(t_i)(S_j(h) \times \{t_j\})} \right) > 0. \end{aligned}$$

We therefore conclude by (D.2) that

$$\sigma_j^{t_i|t_j}(h'')(s_j(h'')) = st \left(\frac{\beta_i(t_i)(S_j(h''), s_j(h'')) \times \{t_j\}}{\beta_i(t_i)(S_j(h'') \times \{t_j\})} \right) > 0$$

for all $h'' \in H_j(s_j)$ weakly following h' . But then, it follows by (D.1) that there is some $\hat{s}_j \in S_j(h')$ with $\hat{s}_j(h'') = s_j(h'')$ for all $h'' \in H_j(s_j)$ weakly following h' such that

$$\sigma_j^{t_i|t_j}|_{h'}(\hat{s}_j) > 0. \tag{D.4}$$

Since t_i is quasi-perfectly rationalizable, we know in particular that t_i induces sequentially rational behavioral strategies. Hence, the induced behavioral strategy $\sigma_j^{t_i|t_j}$ must be sequentially rational for t_j . Since by (D.4) we have that $\sigma_j^{t_i|t_j}|_{h'}(\hat{s}_j) > 0$, it follows that \hat{s}_j must be globally rational for t_j at h' . Since s_j and \hat{s}_j coincide at all $h'' \in H_j(\hat{s}_j)$ that weakly follow h' , it follows that also s_j is




Fig. 4. Procedural quasi-perfect rationalizability does not imply quasi-perfect rationalizability.

globally rational for t_j at h' . But then, we conclude that s_j is globally rational for $b_j^{opp}(t_j)$ at h' . As $h' \in H_j(s_j)$ weakly following h was chosen arbitrarily, it follows that s_j is globally rational for $b_j^{opp}(t_j)$ at every $h' \in H_j(s_j)$ weakly following h . Since we have seen that $b_j^{opp}(t_j) \in B_j^{opp,k-1}$ and $s_j \in S_j^{k-1}(h)$, we conclude that $s_j \in S_j^k(h)$.

We thus see that $st(b_i^{opp}(t_i)(h)(s_j)) > 0$ only if $s_j \in S_j^k$. Since this holds for every $h \in H_i$ and every opponent j , it follows that $b_i^{opp}(t_i)$ believes $S_{-i}^k(h)$ for all $h \in H_i$. As $b_i^{opp}(t_i) \in B_i^{opp,k-1}$, we conclude that $b_i^{opp}(t_i) \in B_i^{opp,k}$, which completes the induction step for (b).

By induction on k , it thus follows that for every player i , every quasi-perfectly rationalizable type $t_i \in T_i$, and every information set $h \in H_i$, (a) every strategy $s_i \in S_i(h)$ that is globally rational for t_i at every $h' \in H_i(s_i)$ weakly following h is in $S_i^k(h)$ for all $k \in \mathbb{N}_0$, and (b) the marginal of $\beta_i(t_i)$ on S_{-i} is in $B_i^{opp,k}$ for all $k \in \mathbb{N}_0$.

Now, take a player i , and a quasi-perfectly rationalizable strategy $s_i \in S_i$. Then, there is an epistemic model $M = (T_i, \beta_i)_{i \in I}$ and a quasi-perfectly rationalizable type $t_i \in T_i$ such that s_i is globally rational for t_i at every $h \in H_i(s_i)$. Then, by (a) above, $s_i \in S_i^k(\emptyset)$ for all $k \in \mathbb{N}_0$, and hence s_i is procedurally quasi-perfectly rationalizable. This completes the proof. \square

We next prove, by means of a counter-example, that the opposite direction of this theorem is not true. Consider the dynamic game in Fig. 4. Note that player 1 is always indifferent between his strategies a and b . We will show that the strategy a is procedurally quasi-perfectly rationalizable, but not quasi-perfectly rationalizable.

Before we give a formal proof, we first provide an informal intuitive argument. According to procedural quasi-perfect rationalizability, player 1 can rationally choose a because he may deem player 2's strategy f infinitely more likely than (d, g) , strategy (d, g) infinitely more likely than (d, h) , and (d, h) infinitely more likely than e . Indeed, under such belief player 1 would assign, at the beginning of the game \emptyset , only non-infinitesimal probability to player 2's strategy f , which is optimal for player 2 from \emptyset onwards if player 2 assigns a high probability to player 1 choosing b .

Such a belief, however, is not possible under the concept of quasi-perfect rationalizability. In order for player 1 to rationally choose a , he must deem player 2's strategy (d, g) at least as likely as (d, h) . Hence, conditional on information set h_2 being reached, player 1 must assign a non-infinitesimal probability to player 2 choosing g . According to quasi-perfect rationalizability, this is only possible if player 1 believes that g is optimal for player 2 at h_2 . Hence, player 1 must believe, conditional on h_2 being reached, that player 2 holds a belief b_2 that assigns probability at least $3/4$ to player 1 having chosen a . Under such a belief b_2 , however, e would be the only optimal strategy for player 2 at h_1 . According to quasi-perfect rationalizability, player 1 must induce a sequentially rational behavioral strategy for player 2. In particular, conditional on player 2's belief b_2 , and conditional on the information set h_1 , player 1 must only assign non-infinitesimal probability to strategies that are optimal for player 2 under the belief b_2 at h_1 . That is, conditional on player 2's belief b_2 , and conditional on the information set h_1 , player 1 must only assign non-infinitesimal probability to strategy e . In particular, this means that player 1 must deem player 2's strategy e infinitely more likely than (d, g) . However, if that is the case player 1's expected utility from choosing a will always be lower than 1, and therefore a cannot be a quasi-perfectly rationalizable strategy.

We will now turn the formal proof. We first show that a survives the procedural quasi-perfect rationalizability procedure. Let $S_i^k(h)$ and $B_i^{opp,k}$ be the sets of strategies and beliefs that survive round k of the procedural quasi-perfect rationalizability procedure, and define $S_i^\infty(h) := \cap_{k \geq 1} S_i^k(h)$ and $B_i^{opp,\infty} := \cap_{k \geq 1} B_i^{opp,k}$. Then, it is easily verified that $S_2^\infty(\emptyset) = \{e, f\}$. Consider player 1's belief

$$b_1^{opp} := (1 - \epsilon - \epsilon^2 - \epsilon^3) \cdot f + \epsilon \cdot (d, g) + \epsilon^2 \cdot (d, h) + \epsilon^3 \cdot e$$

which clearly is in B_1^{opp} . As

$$st(b_1^{opp}(\emptyset)) = f \text{ where } f \in S_2^\infty(\emptyset)$$

it follows that $b_1^{opp}(\emptyset)$ believes $S_{-1}^\infty(\emptyset)$. Hence, $b_1^{opp} \in B_1^{opp,\infty}$.

We now verify that strategy a is globally rational for b_1^{opp} at \emptyset . By construction of the belief b_1^{opp} ,

$$\begin{aligned} u_1(a, b_1^{opp}(\emptyset)) &= u_1(b, b_1^{opp}(\emptyset)) = (1 - \varepsilon - \varepsilon^2 - \varepsilon^3) \cdot 1 + \varepsilon \cdot 2 + \varepsilon^2 \cdot 0 + \varepsilon^3 \cdot 0 \\ &= 1 + \varepsilon - \varepsilon^2 - \varepsilon^3 > 1, \text{ and } u_1(c, b_1^{opp}(\emptyset)) = 1, \end{aligned}$$

and hence a is indeed globally rational for b_1^{opp} at \emptyset . As $b_1^{opp} \in B_1^{opp,\infty}$, it follows that $a \in S_1^\infty(\emptyset)$, and hence a is procedurally quasi-perfectly rationalizable.

We next prove that a is not quasi-perfectly rationalizable. Suppose, contrary to what we want to show, that a were quasi-perfectly rationalizable. Then, there is an epistemic model $M = (T_i, \beta_i)_{i \in I}$ for G , as in Asheim and Perea (2005), and a type $t_1 \in T_1$, such that t_1 is quasi-perfectly rationalizable, and a is globally rational for t_1 at \emptyset . Let $\beta_1(t_1)$ be the cautious non-standard belief that t_1 holds on $S_2 \times T_2$. For every $s_2 \in S_2$, let $\beta_1(t_1)(s_2)$ be the marginal probability that $\beta_1(t_1)$ assigns to s_2 . Then, the expected utility of strategy a at \emptyset for type t_2 is given by

$$u_1(a, t_1, \emptyset) = \beta_1(t_1)(f) \cdot 1 + \beta_1(t_1)(d, g) \cdot 2 + \beta_1(t_1)(d, h) \cdot 0 + \beta_1(t_1)(e) \cdot 0. \tag{D.5}$$

Since $u_1(c, t_1, \emptyset) = 1$ and a is globally rational for t_1 at \emptyset , we must have that $u_1(a, t_1, \emptyset) \geq 1$, which is only possible if $\beta_1(t_1)(d, g) \geq \beta_1(t_1)(d, h)$. Let $t_2 \in T_2$ be such that

$$\beta_1(t_1)((d, g), t_2) \geq \beta_1(t_1)((d, g), t'_2) \text{ for all } t'_2 \in T_2. \tag{D.6}$$

Since $\beta_1(t_1)(d, g) \geq \beta_1(t_1)(d, h)$ we must have, by (D.6), that $\beta_1(t_1)((d, g), t_2)$ is not of infinitely smaller size than $\beta_1(t_1)((d, h), t_2)$. This implies, by (D.2), that $\sigma_2^{t_1|t_2}|_{h_2}(d, g) > 0$. Since t_1 is quasi-perfectly rationalizable, the behavioral strategy $\sigma_2^{t_1|t_2}$ must be sequentially rational for t_2 . In particular, $\sigma_2^{t_1|t_2}|_{h_2}(d, g) > 0$ implies that (d, g) must be globally rational for t_2 at h_2 . This, in turn, is only possible if $\beta_2(t_2)(a) \geq \frac{3}{4}$. Hence, the only strategy that is globally rational for t_2 at h_1 is e . Since the behavioral strategy $\sigma_2^{t_1|t_2}$ must be sequentially rational for t_2 , we must have that $\sigma_2^{t_1|t_2}|_{h_1}(e) = 1$. Hence, in particular, $\beta_1(t_1)(e, t_2)$ must be of infinitely larger size than $\beta_1(t_1)((d, g), t_2)$. But then, it follows by (D.6) that $\beta_1(t_1)(e)$ is of infinitely larger size than $\beta_1(t_1)(d, g)$. However, this insight, together with (D.5), would imply that $u_1(a, t_1, \emptyset) < 1$, and hence a cannot be globally rational for t_1 at \emptyset . That is a contradiction. We thus conclude that a cannot be quasi-perfectly rationalizable. Hence, we have found a strategy a that is procedurally quasi-perfectly rationalizable, but not quasi-perfectly rationalizable.

In fact, we can show even more in this example. The strategy a is not only procedurally quasi-perfectly rationalizable, it is even perfect backwards rationalizable. To see this, let $S_i^k(h)$ and B_i^k be the sets of strategies and beliefs that survive round k of the perfect backwards rationalizability procedure, and define $S_i^\infty(h) := \cap_{k \geq 1} S_i^k(h)$ and $B_i^\infty := \cap_{k \geq 1} B_i^k$. Then, it may be verified that $S_2^\infty(\emptyset) = \{e, f\}$. Consider player 1's belief $b_1 = (b_1^{self}, b_1^{opp})$ where

$$b_1^{self} := \frac{1}{2}(1 - \varepsilon) \cdot a + \frac{1}{2}(1 - \varepsilon) \cdot b + \varepsilon \cdot c$$

and

$$b_1^{opp} := (1 - \varepsilon - \varepsilon^2 - \varepsilon^3) \cdot f + \varepsilon \cdot (d, g) + \varepsilon^2 \cdot (d, h) + \varepsilon^3 \cdot e.$$

Then,

$$\begin{aligned} u_1(a, b_1(\emptyset)) &= u_1(b, b_1(\emptyset)) = (1 - \varepsilon - \varepsilon^2 - \varepsilon^3) \cdot 1 + \varepsilon \cdot 2 + \varepsilon^2 \cdot 0 + \varepsilon^3 \cdot 0 \\ &= 1 + \varepsilon - \varepsilon^2 - \varepsilon^3 > 1, \text{ and } u_1(c, b_1^{opp}(\emptyset)) = 1, \end{aligned}$$

which implies that choices a and b are locally rational for b_1 at \emptyset . Therefore, $S_1^{rat}(b_1, \emptyset) = \{a, b\}$. As $b_1^{self}(\emptyset)$ believes $\{a, b\}$, we conclude that b_1 believes in his own future rationality. Note that player 1 only makes a choice at \emptyset , and therefore he trivially deems his own mistakes least likely under the belief b_1 . We thus conclude that $b_1 \in B_1^0$.

Since

$$st(b_1^{opp}(\emptyset)) = f \text{ where } f \in S_2^\infty(\emptyset)$$

it follows that $b_2^{opp}(\emptyset)$ believes $S_{-1}^\infty(\emptyset)$. It therefore follows that $b_1 = (b_1^{self}, b_1^{opp}) \in B_1^\infty$.

Since we have seen above that a is locally rational for b_1 at \emptyset , it follows that $a \in S_2^\infty(\emptyset)$, and hence a is perfect backwards rationalizable. We have thus found a strategy a that is perfect backwards rationalizable but not quasi-perfectly rationalizable. This means that Theorem 4.1 is no longer true if procedural quasi-perfect rationalizability is replaced by quasi-perfect rationalizability.

Data availability

No data was used for the research described in the article.

References

- Asheim, G.B., Perea, A., 2005. Sequential and quasi-perfect rationalizability in extensive games. *Games Econ. Behav.* 53, 15–42.
- Baltag, A., Smets, S., Zvesper, J.A., 2009. Keep 'hoping' for rationality: a solution to the backward induction paradox. *Synthese* 169, 301–333. (Knowledge, Rationality and Action 705–737).
- Battigalli, P., Siniscalchi, M., 2002. Strong belief and forward induction reasoning. *J. Econ. Theory* 106, 356–391.
- Battigalli, P., Di Tillio, A., Samet, D., 2013. Strategies and interactive beliefs in dynamic games. In: Acemoglu, D., Arellano, M., Dekel, E. (Eds.), *Advances in Economics and Econometrics*. Cambridge University Press, pp. 391–422.
- Battigalli, P., De Vito, N., 2021. Beliefs, plans, and perceived intentions in dynamic games. *J. Econ. Theory* 195. <https://doi.org/10.1016/j.jet.2021.105283>.
- Blume, L.E., Brandenburger, A., Dekel, E., 1991. Lexicographic probabilities and choice under uncertainty. *Econometrica* 59, 61–79.
- Blume, L.E., Meier, M., 2019. Perfect quasi-perfect equilibrium. Downloadable at <https://irihs.ihs.ac.at/id/eprint/4970/1/ihs-working-paper-2019-blume-meier-perfect-quasi-perfect-equilibrium.pdf>.
- Börgers, T., 1994. Weak dominance and approximate common knowledge. *J. Econ. Theory* 64, 265–276.
- Brandenburger, A., 1992. Lexicographic probabilities and iterated admissibility. In: Dasgupta, P., et al. (Eds.), *Economic Analysis of Markets and Games*. MIT Press, Cambridge, MA, pp. 282–290.
- van Damme, E., 1984. A relation between perfect equilibria in extensive form games and proper equilibria in normal form games. *Int. J. Game Theory* 13, 1–13.
- Dekel, E., Fudenberg, D., 1990. Rational behavior with payoff uncertainty. *J. Econ. Theory* 52, 243–267.
- Halpern, J.Y., 2010. Lexicographic probability, conditional probability, and nonstandard probability. *Games Econ. Behav.* 68, 155–179.
- Hammond, P.J., 1994. Elementary non-Archimedean representations of probability for decision theory and games. In: Humphreys, P. (Ed.), *Patrick Suppes: Scientific Philosopher*, vol. 1. Kluwer, Dordrecht, pp. 25–49.
- Penta, A., 2015. Robust dynamic implementation. *J. Econ. Theory* 160, 280–316.
- Perea, A., 2014. Belief in the opponents' future rationality. *Games Econ. Behav.* 83, 231–254.
- Robinson, A., 1973. Function theory on some nonarchimedean fields. *Am. Math. Mon.* 80, 87–109.
- Rubinstein, A., 1991. Comments on the interpretation of game theory. *Econometrica* 59, 909–924.
- Selten, R., 1975. Reexamination of the perfectness concept for equilibrium points in extensive games. *Int. J. Game Theory* 4, 25–55.